

- Hellmayr, C. E. 1913. Critical notes on the types of little known species of neotropical birds. Part II. *Novit. Zool.* 20: 227–256.
- Hellmayr, C. E. 1929. A contribution to the ornithology of northeastern Brazil. *Field Mus. Nat. Hist. Zool. Ser.* 12: 235–501.
- Laubmann, A. 1939. *Die Vogel von Paraguay*. Vol. 1. Verlag von Strecker und Schroder, Stuttgart.
- Lawrence, G. N. 1863. Descriptions of new species of birds of the families Vireonidae and Rallidae. *Proc. Acad. Nat. Sci. Philadelphia* 1863: 106–107.
- Lawrence, G. N. 1864a. Descriptions of new species of birds of the families Tanagridae, Cuculidae, and Trochilidae, with a note on *Panterpe insignis*. *Ann. Lyc. Nat. Hist. New York* 8: 1–6.
- Lawrence, G. N. 1864b. Catalog of birds collected at the Island of Sombrero, W.I., with observations by A. A. Julien. *Ann. Lyc. Nat. Hist. New York* 8: 93–107.
- Penard, F. P. & Penard, A. P. 1910. *De vogels van Guyana*. Vol. 2. Paramaribo.
- Peters, J. L. 1940. *Check-list of Birds of the World*. Vol. 4. Harvard Univ. Press, Cambridge.
- Pinto, O. M. de O. 1966. *Estudo critico e catalogo remissivo das aves do Terrotorio Federal de Roraima*. Inst. Nac. de Pesq. da Amazonia, Manaus.
- Pinto, O. M. de O. 1978. *Novo catalogo das aves do Brasil*. Part 1. Sao Paulo.
- Ridgway, R. 1916. The birds of North and Middle America. Part 7. *Bull US Nat. Mus.* 50, part 7.
- Sclater, P. L. 1862. *Catalogue of a Collection of American Birds*. N. Trubner and Co., London.
- Sclater, P. L. 1870. Further notes on the cuckoos of the genus *Coccyzus*. *Proc. Zool. Soc. London*, 1870: 165–169.
- Sclater, P. L. & Shelley, G. E. 1891. *Catalogue of the Picariae in the collection of the British Museum*. Brit. Mus. (Nat. Hist), London.
- Steinbacher, J. 1962. Beitrage zur Kenntnis der Vogel von Paraguay. *Abh. Senckenb. Naturf. Ges.* 502: 1–106.
- Steullet, A. B. & Deautier, E. A. 1945. *Catalogo sistematico de las aves de la Republica Argentina*. Pt 4: 733–932. Univ. Nac. de la Plata, Buenos Aires.

Address: Richard C. Banks, US Fish and Wildlife Service, National Museum of Natural History, Washington, D.C. 20560.

© British Ornithologists' Club 1988

Weights of some New Caledonian birds

by Charles A. Ross

Received 6 October 1987

In August and September 1986 I spent 24 days in the Southern District of New Caledonia collecting avian anatomical specimens for use in systematic studies and to provide comparative material for identifying collections of Holocene fossils (see Balouet & Olson, in press). Of the 360 specimens obtained, 253 were Passeriformes and included 21 of the 24 native, resident species of passerines (Hannecart & Letocart 1980, 1983).

Collections were made at 5 localities: Mount Dzumac, c. 25 km NE of Noumea; La Foa and vicinity; Bouloupari and vicinity; Sarramea; and the Reserve Forestiere du Col D'Amieu, c. 10 km NE of Sarramea. The Col D'Amieu and Mount Dzumac localities are at 400–1000 m a.s.l. and forested. The other localities are on the western coastal plain and are open savannah with scrub and secondary forest, except the Sarramea locality which is mixed forest and agricultural lands (coffee and fruit trees).

Weights were obtained using pesola spring scales (10, 30, 100, 300 and 1000 g). I know of no published records of weights for many of the following species, although Warner (unpubl. doctoral thesis 1947) presents weights for most species taken from specimens obtained by Thomas L. Macmillan while he was collecting in New Caledonia for the Whitney South Seas Expeditions (1937–1939).

In instances where the sex of specimens is noted, sex was determined by dissection and examination of gonads. Species level nomenclature follows Morony *et al.* (1975). All specimens are deposited in the National Museum of Natural History, Smithsonian Institution, and the Museum National d'Histoire Naturelle, Paris.

The weight of each individual to the nearest 0.5 g (for birds less than 50 g) or 1 g (for heavier birds) is listed for samples of 6 or fewer birds. The mean, standard deviation, range and sample size are given for larger samples. In instances where samples are analysed by sex, the entire sample including unsexed birds is presented first. A query indicates sex unknown.

- Tachybaptus novaehollandiae leucosternos* ♂ 242; ♀ 189
Halietor m. melanoleucos (adult) ♂ 920; ♀ 570, 570, 630, 670, 700, 730; ?680
Nycticorax c. caledonicus ♂ 810
Anas poecilorhyncha pelewensis ♂ 990; ♀ 960
Accipiter haplochrous ♂ 152, 152, 162; ♀ 227, 268, 268
Porphyrio poliocephalus caledonicus ♂ 840, 840; ♀ 690, 690, 820 gravid
Columba vitiensis hypoenochroa ♂ 410, 430
Drepanoptila holosericea ♂ 210; ?220
Ducula goliath ♀ 600
Trichoglossus haematodus deplanchi ♂ 109
Cyanoramphus novaezelandiae saisseti ?72
Collocalia esculenta albidior ♂ 5.0; ♀ 5.0, 5.5
Collocalia spodiopygia leucopygia 6.8 ± 0.6 (6.0–8.5, N=31); ♂ 6.6 ± 0.5 (6.0–7.5, N=10); ♀ 7.1 ± 0.6 (6.0–8.5, N=14)
Halcyon sancta canacorum ♂ 36.5, 41.0, 41.5, 43.0; ♀ 39.5, 43.5
Aplonis s. striata ♂ 53, 60; ?49.5
Artamus leucorhynchus melaleucus ♂ 33.0, 36.0, 37.0; ♀ 31.5, 36.0, 39.5
Corvus moneduloides 275.4 ± 31.5 (230–330, N=22); ♂ 289.3 ± 9.9 (280–310, N=7); ♀ 275.4 ± 36.3 (230–330, N=13)
Coracina analis ♂ 97; ♀ 70
Coracina c. caledonica 138.6 ± 16.0 (117–180, N=21); ♂ 149.1 ± 13.6 (135–180, N=8); ♀ 132.8 ± 17.1 (117–170, N=9)
Lalage leucopygia montrosieri ♂ 16.5, 21.0
Gerygone f. flavolateralis 6.3 ± 0.3 (5.5–6.5, N=12)
Megalururus mariei 25.0 ± 2.7 (20.05–29.25, N=14); ♂ 26.9 ± 1.5 (24.45–29.25, N=8); ♀ 20.05, 22.1, 22.3, 22.7, 23.5, 24.45 (from Warner 1947)
Eopsaltria flaviventris ♂ 14.0; ♀ 10.5, 14.5
Clytorhynchus p. pachycephaloides ♂ 24.5, 25.0
Myiagra c. caledonica 10.8 ± 0.7 (10.0–12.0, N=11)
Rhipidura fuliginosa bulgeri 6.6 ± 0.7 (6.0–8.0, N=11)
Rhipidura spilodera verreauxi ♂ 11.0, 11.2, 12.0; ♀ 9.2, 9.7, 9.85, 10.2 (from Warner 1947)

- Pachycephala caledonica* 20.9 ± 1.3 (18.0–23.0, N=17)
Pachycephala rufiventris xanthetreae 18.2 ± 1.8 (15.5–23.0, N=20); ♂ 18.4 ± 1.5 (16.5–22.0, N=11); ♀ 17.6 ± 1.5 (15.5–19.0, N=7)
Zosterops lateralis griseonota 11.6 ± 0.9 (10.5–13.0, N=7)
Zosterops xanthochroa 10.3 ± 0.9 (8.5–12.0, N=46); ♂ 10.4 ± 0.8 (9.0–12.0, N=18); ♀ 9.9 ± 0.8 (9.0–11.5, N=13)
Gymnomyza aubryana ♂ 211.0, 230.6, 284.2; ♀ 151.8, 159.3 (from Warner 1947)
Lichmera i. incana 13.1 ± 2.0 (9.5–14.5, N=8)
Myzomela sanguinolenta caledonica ♂ 7.5, 8.2; ♀ 6.9, 6.6
Philemon diemenensis 69.3 ± 8.9 (53.0–84.0, N=22); ♂ 74.5 ± 3.9 (69.5–84.0, N=9); ♀ 53.0, 55.0, 56.0, 62.0, 62.0, 62.0
Phylidonyris undulata ♀ 16.0
Erythrura psittacea ♂ 11.5

Acknowledgements

Field work was supported by the Smithsonian Institution Research Opportunities Fund and was facilitated through the cooperation of the Museum National d'Histoire Naturelle, Paris, L'Institut Francaise de Recherche Scientifique pour le Developpement en Cooperation (ORSTOM) and the Service des Forets et du Patrimoine Natural, Noumea. I wish to thank J. C. Balouet and S. Brochot for assistance in the field and for facilitating gun, collection and export permits. S. Tillier (ORSTOM) provided laboratory space in Noumea. I thank S. L. Olson and R. L. Zusi for comments on the manuscript.

References:

- Balouet, J. C. & Olson, S. L. In press. Fossil birds from Holocene deposits in New Caledonia. *Smithsonian Contributions to Paleobiology*.
Hannecart, F. & Letocart, Y. 1980. *Oiseaux de Nouvelle Caledonie et des Loyalties*. Tome 1. Clark & Matheson.
Hannecart, F. & Letocart, Y. 1983. *Oiseaux de Nouvelle Caledonie et des Loyalties*. Tome 2. Dai Nippon, Hong Kong.
Morony, J. L. Jr, Bock, W. J. & Farrand, J. Jr. 1975. *Reference List of the Birds of the World*. American Museum of Natural History.
Warner, D. W. 1947. The Ornithology of New Caledonia and the Loyalty Islands. Unpubl. Doctoral thesis, Cornell University, Ithaca.

Address: Charles A. Ross, Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560 USA.

© British Ornithologists' Club 1988

Variation in the Ground Woodpecker *Geocolaptes olivaceus*

by P. A. Clancey

Received 13 October 1987

The endemic austral African Ground Woodpecker *Geocolaptes olivaceus* (Gmelin), 1788: Cape of Good Hope, is restricted to the Cape Province south of the Orange R., extending north of this in the east of its tenuous



Ross, C A. 1988. "WEIGHTS OF SOME NEW CALEDONIAN BIRDS." *Bulletin of the British Ornithologists' Club* 108, 91–93.

View This Item Online: <https://www.biodiversitylibrary.org/item/123885>

Permalink: <https://www.biodiversitylibrary.org/partpdf/77120>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: British Ornithologists' Club

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.