

TABLE 5. Results of RUMMAGE ANOVA testing occurrence of snails with environmental variables Depth (D), Flow (F) and Slope (S). (N = 262).

Source	df	ss	MS	F	Significance
D	3	31.70	10.58	4.32	0.006
F	1	24.52	24.52	10.02	0.002
S	2	0.72	0.36	0.15	0.863
DF	2 ^a	8.80	4.40	1.80	0.169
DS	6	15.25	2.54	1.04	0.403
FS	2	8.62	4.31	1.76	0.175
Error	141	345.05	2.45		

R² = 0.193
R² adjusted for df = 0.101
^adf lost due to missing cell.

higher, coiled shell. The early literature (Pilsbry 1926, Woodbury 1929, Chamberlain and Jones 1929) states that the purpose of the limpetlike shell and large foot is to allow the snail to live on the vertical surfaces in the trickling water. However, most snails are not found in visibly moving water and are not restricted to vertical surfaces. Slope was not a significant factor (Table 5), and snails are found on horizontal surfaces.

The hypothesis (Woodbury 1933) that the large foot and reduced spire evolved from a more typical physoid type snail as a result of high flows or floods in the seep environment is probably correct. We feel the selective value of attachment during high flows is quite significant. Seeps near the river are occasionally flooded (Malanson 1978), and the ability to remain attached during floods would be an advantage. Based on the experiments, the higher, coiled shell of a typical snail would have caused it to be washed away even in a minor flood, whereas a majority of the Zion Snails would have remained attached to the seep. The adaptive value to the Zion Snail of a large foot and limpetlike shell is, in our opinion, to remain attached during periods of fast flows and flooding, allowing the snail to remain in and exploit seep habitats.

SUMMARY

This research, along with past studies on the Zion Snail, provides a basic understanding of its general distribution and habitat use. Future studies should be done to determine the distribution further up the Narrows and in Orderville Canyon; the role of temperature in controlling reproduction, food habits, and mortality factors; the role of drying seeps on distribution; and, finally, the effect of floods on the Narrows and Orderville Canyon populations. Some of these proposed studies would necessitate collecting

adult snails and raising them in the laboratory, but others could only be accomplished after long-term field observations. Lastly, it must be recognized that we know nothing of populations in gardens high up on the canyon wall or if the snail even exists in those gardens.

Although the snail is endemic to Zion National Park, there are no large populations, but the populations we studied contained sufficient numbers for reproduction to take place. The Zion Snail has probably never existed in large numbers and, in comparison to other snails, it may be considered rare.

LITERATURE CITED

CHAMBERLAIN, R. V., AND D. T. JONES. 1929. A descriptive catalog of the Mollusca of Utah. Bulletin of the University of Utah 19: 1-203.

GREGG, W. O. 1940. Mollusca of Zion National Park, Utah. Nautilus 54: 30-32.

MALANSON, G. P. 1978. Distribution of plant species in the hanging gardens of the Narrows, Zion National Park, Utah. Unpublished thesis, University of Utah, Salt Lake City.

PILSBRY, H. A. 1926. A fresh-water snail, *Physa zionis*, living under unusual conditions. Proceedings of the Academy of Natural Sciences of Philadelphia 77: 325-328.

RYAN, T. A., B. L. JOINER, AND B. F. RYAN. 1981. MINITAB, Minitab reference manual. MINITAB Project, Statistics Department, Pennsylvania State University, University Park, Pennsylvania.

SCOTT, D. T., M. W. CARTER, AND G. R. BRYCE. 1982. RUMMAGE. Statistics Department, Brigham Young University, Provo, Utah.

TALMADGE, R. R. 1970. How far ecology? Of Sea and Shore 1: 131.

WELSH, S. L., AND C. A. TOFT. 1981. Biotic communities of hanging gardens in southeastern Utah. National Geographic Society Research Reports 13: 663-681.

WOODBURY, A. M. 1929. The snails of Zion National Park. Nautilus 43: 54-61.

———. 1933. Biotic relationships of Zion Canyon, Utah, with special reference to succession. Ecological Monographs 3: 147-245.

BLOCKAGE AND RECOVERY OF NITRIFICATION IN SOILS EXPOSED TO ACETYLENE¹

Steven J. Burns^{2,3} and Richard E. Terry²

ABSTRACT.—Acetylene gas is very useful in laboratory and in situ assay procedures for nitrogen fixation and denitrification. There is concern, however, that measurements of denitrification may be underestimated because nitrification, a major source of nitrate, is inhibited by C_2H_2 . The objective of this study was to examine the effects of C_2H_2 partial pressure and length of exposure time on nitrification in soils. Acetylene partial pressures of 0.1, 1.0, and 10.0 kPa were found to effectively inhibit nitrification in soil samples incubated in the laboratory. Both the partial pressure of C_2H_2 and the length of exposure time were found to affect the recovery time of nitrification in soil samples. Nitrification recovered within seven days in samples exposed to 0.1 and 1.0 kPa C_2H_2 for only 24 hours. The recovery of nitrification in samples exposed to 10.0 kPa C_2H_2 for 24 hours or to 0.1 and 1.0 kPa C_2H_2 for 216 hours was delayed for an additional seven days, however.

Acetylene gas has been effectively used in laboratory and in situ techniques for the measurement of both nitrogen fixation and denitrification. The use of acetylene (C_2H_2) is attractive due to the low cost and high sensitivity of the procedures. For N_2 fixation studies, C_2H_2 is used to saturate enzymes responsible for fixation; in the process, C_2H_2 is reduced to ethylene (C_2H_4). Ethylene can be readily detected by gas chromatography and N_2 fixation rates estimated (Hardy et al. 1968, Bergerson 1980). Acetylene has an inhibitory effect on the bacterial enzymes that reduce N_2O to N_2 ; therefore, C_2H_2 has been used in denitrification studies (Balderson et al. 1976, Yoshinari et al. 1977). In the presence of C_2H_2 partial pressures greater than 0.1 kPa, N_2O is the sole gaseous product of denitrification and is detectable by gas chromatography.

Concerns have been expressed about the use of C_2H_2 in denitrification studies, because C_2H_2 has been found to be an effective nitrification inhibitor at a partial pressure of 0.01 kPa (Walter et al. 1979, Berg et al. 1982). In some situations denitrification measurements may be affected by C_2H_2 , because nitrification, a major source of NO_3^- , is inhibited. This is not a problem with short-term laboratory incubations or experiments that involve the addition of nitrate to the soil, but it may be inappropriate to use acetylene in experiments

where NH_4^+ is used as the starting point in denitrification studies or in long-term experiments where mineralization and subsequent nitrification could be expected to produce significant nitrate.

Researchers have studied nitrification in the presence of C_2H_2 partial pressures ranging from 1,000 to 0.01 Pa (Walter et al. 1979, Berg et al. 1982). The minimum effective partial pressure that inhibited nitrification was 10 Pa. However, partial inhibition was found at 0.1 Pa (Berg et al. 1982). These researchers reported that the inhibitory effects of the low C_2H_2 levels (1,000 to 10 Pa) ceased within 7 to 10 days of removal of C_2H_2 .

Acetylene partial pressures of 10 kPa are routinely used in nitrogen fixation studies (Hardy et al. 1968, Bergerson 1980) and in laboratory denitrification studies (Yoshinari et al. 1977, Terry and Tate 1980a, 1980b, Terry et al. 1981). Field measurement of denitrification by the C_2H_2 inhibition technique involves introduction of C_2H_2 to the soil atmosphere through perforated tubing or C_2H_2 treated irrigation water (Ryden, I., *Laboratory evaluation*, 1979; Ryden, II., *Development and application*, 1979, Rolston et al. 1982, Hallmark and Terry 1985). It is likely, with these procedures, that C_2H_2 partial pressures in excess of 10 kPa will exist in portions of the soil atmosphere that will maintain minimum effective levels of 0.1 kPa throughout the soil atmosphere.

¹This publication is based on work supported in part by funds provided by the Bureau of Land Management.

²Department of Agronomy and Horticulture, Brigham Young University, Provo, Utah 84602.

³Present address: Department of Agronomy, University of Missouri, Columbia, Missouri 65201.

TABLE 1. The properties of soils used in this investigation.

Soil series	pH	Sand	Silt	Clay	Organic-C	Total-N
				g kg ⁻¹		
Timpanogos	7.9	348	309	343	12.5	1.27
Woodrow	8.1	239	390	375	8.8	1.40

The effects of C_2H_2 partial pressures greater than 1 kPa on nitrification and the resumption of nitrification following exposure have not been studied. The objective of this research was to determine the effects of various C_2H_2 partial pressures and length of C_2H_2 exposure time on nitrification in soils.

MATERIALS AND METHODS

Soils used in this investigation were Timpanogos clay loam (a fine, loamy, mixed mesic Calcic Argixeroll) collected at the Brigham Young University Agriculture Station near Spanish Fork, Utah, and Woodrow clay loam (a fine silty, mixed mesic Xeric Torrifluvent), collected at Camp Floyd State Park, Fairfield, Utah. The properties of the soils used in this study are presented in Table 1. Air-dried soil samples were analyzed for total N by the micro-Kjeldahl method of Bremner (1965) and for organic C by the method of Allison (1965). The soil pH was measured by glass electrode on a 1:1 soil:water ratio.

Duplicate moist soil samples, the equivalent of 10.0 g dry weight, were placed into 120 ml serum bottles and sealed with septum stoppers. The samples were brought to approximately -0.333 MPa matric potential by addition of 1.25 ml of a solution containing 3.96 g $(NH_4)_2 SO_4 L^{-1}$ to equal an application rate of 100 mg N kg⁻¹ soil. The soils were preincubated for 24 hours before treatment with C_2H_2 .

To compare the inhibitory effects of various levels of C_2H_2 with those of the commercial nitrification inhibitor, nitrapyrin, soil samples were continuously exposed to C_2H_2 partial pressures of 0.1, 1.0 or 10.0 kPa. Impurities were removed from C_2H_2 by passage of the gas through concentrated H_2SO_4 and water traps (Hardy et al. 1968). Nitrapyrin was dissolved in the ammonium solution and added to the soil samples at the rate of 2 mg kg⁻¹. The continuously exposed samples were unsealed, aerated for 10 minutes once a week, then

reexposed to acetylene, thus eliminating anaerobic conditions. Nitrapyrin samples were also aerated for 10 minutes once a week.

To examine the effects of various partial pressures of acetylene on nitrification and the recovery of nitrification following exposure, acetylene was added at rates of 0, 0.1, 1.0, and 10.0 kPa for a period of 24 hours and then removed by flushing the incubation vessels with air. The samples were then incubated at 21 C and continuously aerated with laboratory air at 100% relative humidity using a manifold delivery system and an aquarium air pump. The air flow rate for each sample was 0.5 mL s⁻¹.

To determine the effects of length of acetylene exposure time on nitrification, soil samples were exposed to 0.1, 1.0, or 10.0 kPa C_2H_2 for 24 or 216 hours. Following exposure, acetylene was removed by flushing with air and the samples were then continuously aerated.

Sufficient samples were prepared to allow for duplicate analyses at 0, 7, 14, 21, and 28 days. At the end of incubation inorganic N was extracted from the samples with 50 mL of 2M KCl and NH_4^+ , NO_2^- , and NO_3^- were determined by the steam distillation procedure of Bremner and Keeney (1965).

RESULTS AND DISCUSSION

The effectiveness of C_2H_2 and the commercial nitrification inhibitor, nitrapyrin, on inhibition of nitrification in Timpanogos clay loam was tested. Nitrapyrin, and C_2H_2 at partial pressures ranging from 0.1 to 10.0 kPa, effectively inhibited nitrification in this soil throughout the 28-day incubation (Table 2). The soil samples were preincubated for 1 day to allow the $(NH_4)_2SO_4$ or $(NH_4)_2 SO_4 +$ nitrapyrin solutions to equilibrate with the soil prior to C_2H_2 addition and incubation. The difference in the $NH_4^+ - N$ levels between the nitrapyrin and C_2H_2 treatments at day 0 indicated that approximately 20% of the added

TABLE 2. The effects of various acetylene partial pressures and nitrapyrin on nitrification in Timpanogos clay loam.

C ₂ H ₂	Form	Days of incubation				
Partial pressure #	of N	0	7	14	21	28
kPa		Inorganic N mg kg ⁻¹				
0.1	NH ₄ ⁺	79	79	82	83	81
	NO ₃ ⁻ +NO ₂ ⁻	60	62	60	61	60
1.0	NH ₄ ⁺	80	81	83	84	84
	NO ₃ ⁻ +NO ₂ ⁻	60	61	62	61	62
10.0	NH ₄ ⁺	80	80	81	84	87
	NO ₃ ⁻ +NO ₂ ⁻	59	61	61	59	61
NITRAPYRIN	NH ₄ ⁺	98	95	97	97	n.d.*
	NO ₃ ⁻ +NO ₂ ⁻	55	51	56	53	n.d.*

*n.d. = not determined
#Except Nitrapyrin

TABLE 3. The effects of a 24-hour exposure to C₂H₂ partial pressures ranging from 0 to 10 kPa on nitrification in Timpanogos cl.

C ₂ H ₂	Form	Days of incubation				
Partial pressure	of N	0	7	14	21	28
kPa		Inorganic N mg kg ⁻¹				
0	NH ₄ ⁺	52	0	0	0	0
	NO ₃ ⁻ +NO ₂ ⁻	83	140	147	148	150
0.1	NH ₄ ⁺	78	0	0	0	0
	NO ₃ ⁻ +NO ₂ ⁻	61	146	146	152	148
1.0	NH ₄ ⁺	70	0	0	0	0
	NO ₃ ⁻ +NO ₂ ⁻	63	144	144	152	156
10.0	NH ₄ ⁺	68	45	2	0	0
	NO ₃ ⁻ +NO ₂ ⁻	63	93	140	150	155

NH₄⁺-N was nitrified during preincubation. The finding that C₂H₂ partial pressures ranging from 0.1 to 10.0 kPa inhibited nitrification concurred with earlier work of Walter et al. (1979) and Berg et al. (1982), who showed that C₂H₂ partial pressures of 1.0 and 0.1 kPa effectively inhibited nitrification in soils.

The effects of 24 hours of exposure to C₂H₂ partial pressures ranging from 0 to 10.0 kPa on subsequent changes in NH₄⁺-N and (NO₃⁻ + NO₂⁻)-N concentrations in the Timpanogos soil are shown in Table 3. The zero time of incubation followed the 24-hour preincubation and the 24 hours of exposure of C₂H₂. Nitrification continued in the control samples (0 kPa) during the 24-hour period that the remaining treatments were exposed to C₂H₂. For this reason more (NO₃⁻ + NO₂⁻)-N had accumulated in the control samples during preincubation. During the first week if incubation following removal of C₂H₂, nitrification of added NH₄⁺-N was complete in all samples except those treated with 10.0 kPa. Nitrification in the samples exposed to 10.0 kPa C₂H₂ for 24 hours was slowed for approxi-

mately two weeks due to the lingering effects of acetylene exposure. There were no differences in the NH₄⁺-N and (NO₃⁻+NO₂⁻)-N concentrations during incubation of the 0.1 and 1.0 kPa treatments. During the first week following removal of C₂H₂ from the samples, the nitrification rate in samples exposed to 0.1 and 1.0 kPa C₂H₂ was 12 mg N kg⁻¹ day⁻¹ compared to 4.3 mg N kg⁻¹ day⁻¹ in those treated with 10 kPa.

Samples of the Woodrow clay loam were incubated aerobically following a 24-hour exposure to, and subsequent removal of, C₂H₂ partial pressures ranging from 0 to 10.0 kPa. The effects of this brief exposure on subsequent changes in NH₄⁺-N and (NO₃⁻+NO₂⁻)-N concentrations are shown in Table 4. During the first week following C₂H₂ removal, the nitrification rates for 0, 0.1, 1.0, and 10.0 kPa C₂H₂ treatments were 5.2, 5.7, 2.1, and 3.6 mg N kg⁻¹ day⁻¹, respectively. Accumulation of (NO₃⁻+NO₂⁻)-N in the samples exposed to 1.0 and 10.0 kPa was slower during the first week of incubation.

Nitrification proceeded in the Woodrow cl, an unfertilized rangeland soil, at a slower pace

TABLE 4. The effects of a 24-hour exposure of C₂H₂ partial pressures ranging from 0 to 10 kPa on nitrification in Woodrow cl.

C ₂ H ₂	Form	Days of incubation				
Partial pressure	of N	0	7	14	21	28
kPa		Inorganic N mg kg ⁻¹				
0	NH ₄ ⁺	79	31	0	0	0
	NO ₃ ⁻ +NO ₂ ⁻	19	56	105	106	114
0.1	NH ₄ ⁺	87	54	5	0	0
	NO ₃ ⁻ +NO ₂ ⁻	12	52	98	104	111
1.0	NH ₄ ⁺	83	70	19	0	0
	NO ₃ ⁻ +NO ₂ ⁻	12	27	82	101	111
10.0	NH ₄ ⁺	83	62	18	2	0
	NO ₃ ⁻ +NO ₂ ⁻	13	38	83	93	110

TABLE 5. The effects of C₂H₂ partial pressure and length of exposure time on nitrification in Timpanogos cl.

C ₂ H ₂	Exposure	Form	Days of incubation				
Partial pressure	time	of N	0	7	14	21	28
kPa	Hours		Inorganic N mg kg ⁻¹				
10	24	NH ₄ ⁺	68	45	2	0	0
		NO ₃ ⁻ +NO ₂ ⁻	63	93	140	151	155
	216	NH ₄ ⁺	81	48	1	0	0
		NO ₃ ⁻ +NO ₂ ⁻	59	89	149	149	148
1.0	24	NH ₄ ⁺	70	0	0	0	0
		NO ₃ ⁻ +NO ₂ ⁻	63	144	145	153	156
	216	NH ₄ ⁺	81	47	1	0	0
		NO ₃ ⁻ +NO ₂ ⁻	60	95	139	148	158
0.1	24	NH ₄ ⁺	78	0	0	0	0
		NO ₃ ⁻ +NO ₂ ⁻	61	146	146	153	148
	216	NH ₄ ⁺	80	47	1	0	0
		NO ₃ ⁻ +NO ₂ ⁻	61	94	148	152	158

than in the Timpanogos cl, a fertilized cropped soil. The results reported above indicate that brief exposure (24 hours) to C₂H₂ partial pressures ranging from 0.1 to 1.0 kPa have little effect on nitrification in these soils. Brief exposure of soil to 10.0 kPa C₂H₂ slowed subsequent nitrification for as long as two weeks, however.

The effects of length of C₂H₂ exposure time on recovery of nitrification in Timpanogos cl are shown in Table 5. Nitrification rates in soil samples treated with 10.0 kPa acetylene were equivalent during the first week of incubation following C₂H₂ removal whether exposed to the gas for 24 or 216 hours. Nitrification was nearly complete within one week in samples exposed to 0.1 or 1.0 kPa for only 24 hours. Nitrification rates in samples exposed to 0.1 and 1.0 kPa C₂H₂ for 216 hours were much slower, however. The effects of lengthy exposure (216 hours) to C₂H₂ partial pressures of 0.1 and 1.0 kPa on subsequent nitrification were similar to the effects of brief exposure (24 hours) to 10.0 kPa.

Walter et al. (1979) reported that nitrification rates in soil samples exposed to C₂H₂ partial pressures ranging from 0.1 to 1.0 kPa for 24 hours returned to those of control samples after an 8 to 10 day lag period. Similar results were reported by Berg et al. (1982), who exposed soil samples to 0.01 kPa C₂H₂ for seven days. They reported that rates of nitrate production were similar to the rates in control samples seven days after C₂H₂ removal.

The findings of this experiment indicate that both the partial pressure of C₂H₂ and the length of exposure time affect the recovery time of nitrification in soil samples. Exposure of Woodrow cl to 0.1 and 1.0 kPa C₂H₂ for 24 hours slowed nitrification for approximately seven days. Exposure of Timpanogos cl to 0.1 and 1.0 kPa C₂H₂ for 24 hours had little effect on nitrification once the inhibitor was removed. Exposure of this soil to 0.1 and 1.0 kPa acetylene for 216 hours slowed nitrification for approximately seven days, however. These levels of C₂H₂ exposure are commonly used in

laboratory and in situ denitrification studies (Yoshinari et al. 1977, Ryden, *II.*, *Development and application*, 1979; Rolson et al. 1982, Ryden and Dawson 1982). The recovery of nitrification in Timpanogos soil exposed to 10.0 kPa C_2H_2 was delayed for at least seven days whether the samples were exposed to the gas for 24 or 216 hours. Acetylene partial pressures of 10.0 kPa have been used in denitrification studies and in studies of concurrent denitrification and nitrogen fixation (Terry and Tate 1980a, 1980b, Yoshinari et al. 1977).

Problems with the use of C_2H_2 in denitrification studies would likely be encountered in soils where the nitrate supply is limited by nitrification. Ryden (1982) reported that denitrification was underestimated in soil samples incubated in the laboratory in the presence of C_2H_2 for 168 hours. Nitrate became exhausted in the samples during incubation. Problems with nitrification inhibition in denitrification studies may be avoided in laboratory studies by adding supplemental nitrate and/or adopting short incubation periods (<24 hours) (Terry and Tate, 1980b). In the design of field studies on denitrification, it would be wise to rotate study sites every 7 to 14 days to allow nitrification to proceed in the soil. The use of sites previously exposed to C_2H_2 should be avoided for 14 to 21 days because of the slow recovery of nitrification following prolonged exposure to C_2H_2 .

LITERATURE CITED

- ALLISON, F. E. 1965. Organic carbon. In C. A. Black, ed., *Methods soil analysis. Part 2—Chemical and microbiological methods*. Amer. Soc. of Agron., Madison, Wisconsin. Agronomy 9: 1367–1378.
- BALDERSON, W. L., B. SHERR, AND W. J. PAYNE. 1976. Blockage by acetylene of nitrous oxide reduction *Pseudomonas Perfectomarinus*. Appl. Environ. Microbiol. 31: 504–508.
- BERG, P., L. KLEMEDTSSON, AND T. ROSSWALL. 1982. Inhibitory effect of low partial pressure of acetylene on nitrification. Soil Biol. Biochem. 14: 301–303.
- BERGERSON, F. J. 1980. *Methods of evaluating biological nitrogen fixation*. John Wiley and Sons Inc., New York.
- BREMNER, J. M. 1965. Total nitrogen. In C. A. Black, ed., *Methods of soil analysis. Part 2—Chemical and microbiological methods*. Amer. Soc. of Agron., Madison, Wisconsin. Agronomy 9: 1149–1178.
- BREMNER, J. M., AND D. R. KEENEY. 1965. Steam distillation methods for determination of ammonium, nitrate, and nitrite. Anal. Chim. Acta. 32: 485–495.
- HALLMARK, S. L., AND R. E. TERRY. 1985. Field measurement of denitrification in irrigated soils. Soil Sci. 140: 35–44.
- HARDY, R. W. F., R. D. HOLSTEN, E. K. JACKSON, AND R. C. BURNS. 1968. The acetylene-ethylene assay for N_2 fixation: Laboratory and field evaluation. Plant Physiol. 42: 1185–1207.
- ROLSTON, D. E., A. N. SHARPLEY, D. W. TOY, AND F. E. BROADBENT. 1982. Field measurement of denitrification: III. Rates during irrigation cycles. Soil Sci. Soc. of Amer. J. 46: 289–296.
- RYDEN, J. C. 1982. Effects of acetylene on nitrification and denitrification in two soils during incubation with ammonium nitrate. J. Soil Sci. 33: 263–270.
- RYDEN, J. C., AND K. P. DAWSON. 1982. Evaluation of the acetylene-inhibition technique for the measurement of denitrification in grassland soils. J. Sci. Food Agric. 33: 1197–1206.
- RYDEN, J. C., L. J. LUND, AND D. D. FOCHT. 1979. Direct measurement of denitrification loss from soils: I. Laboratory evaluation of acetylene inhibition of nitrous oxide reduction. Soil Sci. Soc. Amer. J. 43: 110–140.
- RYDEN, J. C., L. J. LUND, J. LETEY, AND D. D. FOCHT. 1979. Direct measurement of denitrification loss from soils: II. Development and application of field methods. Soil Sci. Soc. of Amer. J. 43: 110–118.
- TERRY, R. E., AND R. L. TATE. 1980a. Denitrification as a pathway for nitrate removal from organic soils. Soil Sci. 129: 162–166.
- . 1980b. The effect of nitrate on nitrous oxide reduction in organic soils and sediments. Soil Sci. Soc. Amer. J. 44: 744–746.
- TERRY, R. E., R. L. TATE, AND J. M. DUXBURY. 1981. The effect of flooding on nitrous oxide emissions from an organic soil. Soil Sci. 132: 228–232.
- WALTER, H. M., D. R. KEENEY, AND I. R. FILLERY. 1979. Inhibition of nitrification by acetylene. Soil Sci. Soc. Amer. J. 43: 195–196.
- YOSHINARI, T., R. HYNES, AND R. KNOWLES. 1977. Acetylene inhibition of nitrous oxide reduction and measurement of denitrification and nitrogen fixation in soil. Soil Biol. Biochem. 9: 177–183.

NEW THAGRIINE LEAFHOPPERS FROM THE ORIENTAL REGION, WITH A
KEY TO 30 SPECIES (HOMOPTERA: CICADELLIDAE: COELIDIINAE)

M. W. Nielson

ABSTRACT.—Five new species of *Thagria* from the Oriental region are described and illustrated. These include *melichari* from Thailand, *unidentata* from Indonesia, *marissae* from southern China, *bifida* from Nepal, and *insolentis* from an undetermined locality in the Oriental region. There are presently 166 species in this large and unique genus. A key to males of 30 species is included.

The genus *Thagria* Melichar is the largest group of coelidiine leafhoppers. Although they occur primarily in the Oriental region, many species are found in the Australian region (not known in Australia proper) and several are in the southern Palearctic region (southern China, southern Korea, and southern Japan). Prior to 1977 only 36 species were known. Since then 125 species have been described (Kwon and Lee 1979, Nielson 1977, 1980a, 1980b, 1980c, 1980d, 1982). The five new species described herein bring the present total to 166 species.

The genus is uniquely characterized by the males possessing a distinctive and highly diverse ventral paraphysis on which a tubular aeadeagal shaft is attached basally to and freely articulates dorsally with the paraphysis. The many configurations of the ventral paraphysis in combination with highly modified structures of the 10th segment and caudodorsal processes of the pygofer differentiate the numerous species.

A key to males of 30 species including those described in previous papers (except Kwon and Lee 1979) after my 1977 revision and those treated herein is presented. A regional key for all known species will be presented later.

Host plants and biology of species in the group are very poorly known.

Key to Males of *Thagria*

- 1. Clypellus broad, swollen basally or nearly so, basal width equal to or greater than basal width of clypeus, lateral margins usually narrowed medially 2

- Clypellus narrow, never swollen basally, basal width narrower than basal width of clypeus, lateral margins usually parallel, sometimes expanded distally 16
- 2(1). Ventral paraphysis curved ventrally at distal 1/2 to 1/3 in lateral view, apex decurved 3
- Ventral paraphysis not as above, in lateral view straight or recurved 5
- 3(2). Style with apex bifurcate or divided into 2 slender rami 4
- Style not as above (Fig. 16, Nielson 1980a) *blockeri* Nielson
- 4(3). Aedeagus long, extending beyond midlength of ventral paraphysis; 10th segment process with dentate process on middle of dorsal margin (Fig. 2) *bifida*, n. sp.
- Aedeagus shorter, reaching to about midlength of ventral paraphysis; 10th segment process with longer process on ventral margin (Fig. 25, Nielson 1980b) *thailandensis* Nielson
- 5(3). Ventral paraphysis symmetrical 6
- Ventral paraphysis asymmetrical 10
- 6(5). Tenth segment with paired processes 7
- Tenth segment without paired processes (Fig. 13, Nielson 1980b) *ampla* Nielson
- 7(6). Ventral paraphysis without basal paired processes on dorsal margin 8
- Ventral paraphysis with basal paired processes on dorsal margin (Fig. 22, Nielson 1980b) *serrastyla* Nielson
- 8(7). Style very long, exceeding midlength of ventral paraphysis; ventral paraphysis without spines distally 9
- Style very short, not reaching midlength of ventral paraphysis; ventral paraphysis with lateral spines distally (Fig. 28, Nielson 1982) *barbata* Nielson
- 9(8). Style attenuated distally (Fig. 2, Nielson 1982) *fossiata* Nielson

¹Monte L. Bean Life Science Museum, Brigham Young University, Provo, Utah 84602.

— Style forked distally (Fig. 11, Nielson 1980c) *furculata* Nielson

10(5). Ventral paraphysis with basal processes on dorsal margin 11

— Ventral paraphysis without basal processes but with medial or subapical processes on dorsal margin 14

11(10). Ventral paraphysis with paired basal processes 12

— Ventral paraphysis with single basal process. 13

12(11). Basal processes on paraphysis symmetrical (Figs. 11,12) *melichari*, n. sp.

— Basal processes on paraphysis asymmetrical (Fig. 40, Nielson 1982) *hollowayi* Nielson

13(11). Style broad throughout in dorsal view, without dentate subapical processes (Fig. 39, Nielson 1980b) *boulardi* Nielson

— Style narrowed at distal 1/4 in lateral view, with dentate subapical process (Fig. 35, Nielson 1980b) *paraornata* Nielson

14(10). Tenth segment and caudodorsal margin of pygofer with processes of equal length in lateral view; ventral paraphysis with short lateral process distad of middle. 15

— Tenth segment and caudodorsal margin of pygofer with processes of unequal length in lateral view; ventral paraphysis with short lateral process on middle (Fig. 4, Nielson 1980b). *undulata* Nielson

15(14). Tenth segment processes very narrow and sinuate, nearly needlelike at distal 2/3 in dorsal view (Fig. 8, Nielson 1980b) . *capilla* Nielson

— Tenth segment processes broader and nearly straight, not needlelike in dorsal view (Fig. 20, Nielson 1980a) *paradigitata* Nielson

16(2). Ventral paraphysis symmetrical 17

— Ventral paraphysis asymmetrical 25

17(16). Ventral paraphysis keeled ventrally. 18

— Ventral paraphysis not as above 20

18(17). Ventral paraphysis with subbasal ventral keel 19

— Ventral paraphysis with subapical ventral keel (Fig. 10, Nielson 1980d) . *paraloae* Nielson

19(18). Style very long, extending beyond apex of ventral paraphysis (Fig. 22, Nielson 1980d) *samuelsoni* Nielson

— Style very short, extending only to base of ventral paraphysis (Fig. 3, Nielson 1980c) *ventrocarina* Nielson

20(17). Ventral paraphysis with paired basal process on dorsal margin 21

— Ventral paraphysis not as above 22

21(20). Paired basal processes of paraphysis very long, nearly reaching to apex of paraphysis (Fig. 3, Nielson 1980d) *bilateralis* Nielson

— Paired basal processes of paraphysis shorter, not reaching midlength of paraphysis (Figs. 16, 17) *insolentis*, n. sp.

22(20). Caudovernal lobe of pygofer without spines 23

— Caudovernal lobe of pygofer with 2 short spines apically (Fig. 7, Nielson 1982) *bidentata* Nielson

23(22). Ventral paraphysis without lateral processes 24

— Ventral paraphysis with lateral processes subapically (Fig. 3, Nielson 1980a) *srilankensis* Nielson

24(23). Style with subapical bifurcation (Fig. 21, Nielson 1982) *bifurcata* Nielson

— Style without subapical bifurcation (Fig. 11, Nielson 1980a) *brincki* Nielson

25(16). Style with distal half straight or nearly so ... 26

— Style with distal half hooked (Fig. 17, Nielson 1980d) *paraexilis* Nielson

26(25). Ventral paraphysis without ventral keel 27

— Ventral paraphysis with ventral keel subbasally (Fig. 17, Nielson 1982) *mutabilis* Nielson

27(26). Ventral paraphysis with 1–2 lateral processes on or near apex. 28

— Ventral paraphysis without such processes .. 29

28(27). Ventral paraphysis with a pair of unequal distal processes (Fig. 23) *marrisae*, n. sp.

— Ventral paraphysis with a single, large, retrorse lateral process subapically (Fig. 35, Nielson 1982) *retrorsa* Nielson

29(27). Caudovernal lobe of pygofer with a single long spine (Fig. 26) *unidentata*, n. sp.

— Caudovernal lobe of pygofer without such spine (Fig. 13, Nielson 1980c) *kaloostiani* Nielson

Thagria bifida, n. sp.

Figs. 1–6

LENGTH: Male 6.90 mm.

Moderate-sized, slender species. General color black with tannish translucent costa, face black.

Head small, subconical, much narrower than pronotum; crown broad, width about equal to width of eyes, produced beyond anterior margin of eyes, elevated above level of eyes, lateral margins convergent basally; eyes moderately large, semiglobular; pronotum with length about equal to length of crown; scutellum large; forewings long and narrow, venation typical of genus; clypeus long and broad, lateral margins excised near middle; clypellus short and broad, base broad and swollen, lateral margins below converging to truncate apex.



1986. "New thagriine leafhoppers from the Oriental region, with a key to 30 species (Homoptera: Cicadellidae: Coelidiinae)." *The Great Basin naturalist* 46, 321–328. <https://doi.org/10.5962/bhl.part.7734>.

View This Item Online: <https://www.biodiversitylibrary.org/item/33077>

DOI: <https://doi.org/10.5962/bhl.part.7734>

Permalink: <https://www.biodiversitylibrary.org/partpdf/7734>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Brigham Young University

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.