

"6. July 12, 1894. Seven eggs; incubation advanced. Nest in tussock of coarse grass, made of dead grass and bull-rush leaves.

"7. August, 10, 1898. Six eggs; small embryos."—H. H. & C. S. Brimley.—WITMER STONE, *Academy of Natural Sciences, Philadelphia, Pa.*

Breeding of the Little Black Rail (*Porzana jamaicensis*) in New Jersey in 1844 and 1845.—Soon after the above note was written, by a curious coincident I came across a definite account of the breeding of this bird in New Jersey. Apart from a bare statement of the fact in Turnbull's 'Birds of East Pennsylvania and New Jersey,' the record seems not to have been published.

The facts are contained in letters written by Chas. C. Ashmead of Philadelphia to Prof. Baird, and for the privilege of publishing them I am indebted to Miss Lucy H. Baird. The extracts relative to the Black Rail are as follows:

July 28, 1844.—"My brother-in-law has just arrived from the seashore. Not long since he found a nest of the Black Rail; it was on a *fresh* water meadow near the seaside, and contained four eggs. He also caught the bird."

Oct. 2, 1844.—"I have the eggs of the Black Rail, also the full-plumaged male bird, in my possession."

Nov. 17, 1845.—"Tom Beesley has found another nest of the Black Rail, making the third he has found. The two first nests he found, one early in June, 1844, with four eggs; one early in June, 1845, with three eggs; and the last one about the middle of August, with but one egg in the nest. He had killed the bird before he found the nest. All three of the nests were found on the same spot of ground,—a fresh marsh on the banks of the Great Egg Harbor River, and not more than one fourth as big as the College Campus [at Carlisle]."

The locality was evidently Beesley's Point, and a brief mention of the spotting on the bird leaves no doubt of its identity. The fact, however, that Mr. Ashmead and his brother were constantly at the Academy at this time, and in daily association with Mr. Cassin, would preclude the possibility of any error in identification. The discovery that this obscure little bird still breeds in this locality is well within the limits of possibility.—WITMER STONE, *Academy of Natural Sciences, Philadelphia, Pa.*

Occurrence of *Tringa fuscicollis* in Virginia in Autumn.—During a visit to the Eastern Shore of Virginia in company with Dr. William C. Rives, Sept. 21–28, 1899, a specimen of the White-rumped Sandpiper was secured near Chincoteague, Va., on the sand flats lying inside the beach of Assateague Island on Sept. 24. The bird was shot as it flew past with a flock of Semipalmated Sandpipers, among which it was conspicuous by reason of its larger size. It was the only individual of this species observed on the trip. This species appears in Dr. Rives's 'Birds of the



Stone, Witmer. 1900. "Breeding of the Little Black Rail (*Porzana jamaicensis*) in New Jersey in 1844 and 1845." *The Auk* 17, 172-172.

<https://doi.org/10.2307/4069169>.

View This Item Online: <https://www.biodiversitylibrary.org/item/54122>

DOI: <https://doi.org/10.2307/4069169>

Permalink: <https://www.biodiversitylibrary.org/partpdf/89688>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.