

ANNUAL ENERGY BUDGETS FOR THREE COMMON RODENT SPECIES IN THE NORTHERN GREAT BASIN¹

R. Kent Schreiber^{2,3}

ABSTRACT.— Annual energy budgets were calculated for three species of small mammals (*Peromyscus maniculatus*, *Onychomys leucogaster*, *Reithrodontomys megalotis*) from the northern Great Basin, Benton County, Washington. Individual ingestion rates were based on species activity, microclimate regime, coefficient of digestibility, caloric diet, and the cost for reproduction. For males and females, the estimated energy expenditures were: *P. maniculatus*, 6080, 5891; *O. leucogaster*, 5714, 6587; and *R. megalotis*, 4057, 3791 kcal/yr. By comparison, each species on an individual basis processes more energy annually than the more abundant species in the community, *Perognathus parvus*, but their total contribution to community energy flow is apparently minor. Integration of these results with other ecological parameters is necessary to develop new hypotheses on the role of small mammal consumers in cold desert ecosystems.

Historically, ecologists have studied and compared ecosystems and their component species in terms of density and biomass. However, this approach does not emphasize the impact of each species on the total system or its relationship to other trophic levels within the system. The concept of energy flow provides such a common factor for comparing ecosystems and also for evaluating the relative importance and success of the constituent populations.

In the northern part of the Great Basin common rodent species include the Great Basin pocket mouse (*Perognathus parvus*), deer mouse (*Peromyscus maniculatus*), northern grasshopper mouse (*Onychomys leucogaster*) and the western harvest mouse (*Reithrodontomys megalotis*). These small mammal consumers are representative of an important pathway for energy transfer in a cold desert ecosystem. The bioenergetics of the predominant species, *P. parvus*, has been discussed in a previous paper (Schreiber 1978b). This paper reports on the energy budgets of the three, less abundant, species.

Energy flow through a rodent population can be determined from daily energy requirements and ingestion rates of individuals during each season. In this study I calculated ingestion rates by considering activity of each

species in field-encountered microclimates and their resulting metabolic demands. Adjustments in the ingestion rates were made for the additional energy cost of reproduction and for the energy savings while residing in a nest.

MATERIALS AND METHODS

Study Area

The study area is in the Hanford Works Department of Energy (DOE) Reservation 19 km northwest of Richland, Benton County, Washington. The 58-year average annual precipitation for the Reservation is 159 mm. Other climatological and edaphic conditions for the general area have been summarized by Stone, Jenne, and Thorp (1972). Vegetation is mostly typical of the *Artemisia tridentata*-*Poa* association (Daubenmire 1970) with the exception of native grass species. Cheatgrass (*Bromus tectorum*) was introduced into the area over a half century ago and now has replaced native species as the dominant ground cover. Shrubby species present include big sagebrush (*Artemisia tridentata*), bitterbrush (*Purshia tridentata*), and two rabbitbrush species (*Chrysothamnus nauseosus*; *C. viscidiflorus*).

¹This study was supported in part by the U.S.-IBP Desert Biome Program (Grant GB 15886 from the National Science Foundation).

²Department of Biological Sciences, University of Idaho, Moscow, Idaho 83843.

³Present address: U.S. Fish and Wildlife Service, National Power Plant Team, 2929 Plymouth Road, Ann Arbor, Michigan 48105.

Trapping

Although no attempt was made in this study to delineate absolute population numbers, monthly sampling with snap-traps determined species composition, relative population sizes, and trends. Traps were spaced approximately 3 m apart, with 50 traps per 150 m line. Rolled oats paste was used for bait and traps were normally set for three consecutive nights in each trapping session. Sex, weight, and reproductive status of all captures were recorded. In addition, rodents were live-trapped in peripheral areas for use in laboratory food trials.

Microclimate

To establish the microclimate regime of the species, temperatures were recorded continuously at the surface and at a burrow depth of 0.5 m by a seven-day, two-pen thermograph. Data were summarized to coincide with monthly trapping sessions. Mean diurnal surface temperature (T_d) was calculated as the average of even-hour temperatures from dawn to dusk. Correspondingly, mean nocturnal surface temperatures (T_n) were calculated as the average of even-hour temperatures from dusk to dawn. Burrow temperature (T_b) was calculated as the mean of the daily maximum and minimum subsurface temperature.

Digestibility

Energy content of ingested materials was determined by combustion in a semimicro oxygen bomb calorimeter. The coefficient of digestibility (digested proportion of ingested food) was measured directly in the laboratory from food intake and indirectly by the ash-tracer method for free-living animals (Schreiber 1979).

Energy Expenditure

Annual ingestion rates were calculated from the activity and resting time of the species in field-encountered microclimates (surface and burrows) and their resulting caloric demands. The additional energy cost of re-

production and the energy savings from insulating properties of the nest were incorporated into the calculations.

Ingestion rates were calculated using the general model:

$$I = [(E_r + E_a) + E_{g_3}] D^{-1} \\ = (E_m + E_{g_3}) D^{-1} \quad (1)$$

where I is ingestion rate (kcal/yr), E_r and E_a are energy costs during rest and during activity, E_m is their sum (maintenance), E_{g_3} is energy costs for growth from weaning to subadult, and D is coefficient of digestibility. Additional growth between the subadult and adult stage was considered by using the average adult weight when calculating maintenance energy (E_m). During pregnancy and lactation, females incur additional energy demands due to respiration and growth of the embryos. To account for embryonic respiration, gravid females were included in calculations of female mean weight. Ingestion rates for females were calculated as—

$$I_f = (E_m + p E_{g_1} + w E_{g_2} + E_{g_3}) (D^{-1}) \quad (2)$$

—where the coefficient p is mean brood size at parturition (i.e., mean litter size \times average number of litters per year), w is the mean brood size at weaning (i.e., mean brood size (p) minus mortality during nursing period), and E_{g_1} and E_{g_2} are energy costs for growth from conception to birth and from birth to weaning, respectively. Intrauterine mortality is unknown but probably small and has been ignored in the calculations. Females with either embryos or placental scars were recorded as bearing one litter; females with both embryos and scars or scars of an undetermined number were recorded as having two litters. For all species I assumed a conservative survival rate of 80 percent for nursing young (Kaczmarski 1966).

Energy Costs at Rest: Resting metabolic rates (RMR) for individual species were taken from the literature. Since animals in a burrow and occupying a nest have lower energy requirements during rest, I adjusted RMR's for this energy conservation by plotting the nesting metabolic rate (NMR) as a regression line based on 0.81 RMR at 1 C and 0.87 RMR at 12 C (based on data reported for the harvest mouse, Pearson 1960).

Energy Cost of Activity: Metabolic rates increase during periods of activity. Estimates of daily and seasonal variations in amounts of activity were based on field-monitored activity of free-roaming mice tagged with a radioactive nuclide (Schreiber 1973), personal communications from other field investigators, and published data (French et al., 1966). The incremental energy demand (E_a) during these activity period was calculated from information cited in Chew and Chew (1970).

Energy Cost of Growth: Energy cost for growth during a specific stage of development (E_g) is the product of the weight gain (W) and tissue caloric value (K) divided by growth efficiency (G) during that period of growth, i.e.,

$$E_g = (WK)(G^{-1}) \quad (3)$$

I used the following caloric values for the tissues (K): for the embryo, 0.98 kcal/g fresh weight, based on the average caloric values of five species of newborn rodents (Gorecki 1965, Myrcha and Walkowa 1968, Soholt 1973), and for the unweaned young, 1.39 kcal/g, assuming an average weaning age of 25 days and the mean caloric value of two species of rodents (Myrcha and Walkowa 1968, Soholt 1973). Caloric values for weaned young of individual species are given in the results. Growth efficiencies were taken from the literature and based on average values, $G = 13.8$ percent for embryos, 15.0 percent for unweaned young, and 5.0 percent for weaned young (Kaczmarek 1966, Migula 1969, Drozd et al. 1972).

RESULTS AND DISCUSSION

Composition and Abundance

Snap traps effectively sample small mammal populations (Wiener and Smith 1972), and they are particularly applicable for censusing large areas. Under ideal conditions the total number of individuals caught in traps is proportional to population density and reflects the structure of the population (Hansson 1967, Petticrew and Sadlier 1970). Although trapping percentages are not direct estimates of density, they are nonetheless in-

dicative of population trends and therefore provide insight into the influence a particular species has in the transfer of energy into the community.

Trap lines were operated one night each in September and November 1969 and March, April, and May 1970 and at least three nights per month from June 1970 to May 1971 (Table 1). A total of 1470 rodents was captured in 14,289 trap-days (one trap set for one day). The average monthly effort was 794 trap-days and the overall trapping success was 10.3 percent. Field observations indicated traps placed in dense cover or concealed by shadows had somewhat greater success. On other areas of the Hanford Reservation general trapping success has been reported as low as 4 percent on fire-disturbed grasslands with stony soils (Hedlund et al. 1975) and as high as 44 percent in shrub-steppe habitats with coarse-textured sands (O'Farrell 1975b).

A total of four species of rodents was snap-trapped on the study area: the Great Basin pocket mouse, *Perognathus parvus*; deer mouse, *Peromyscus maniculatus*; northern grasshopper mouse, *Onychomys leucogaster*; and western harvest mouse, *Reithrodontomys megalotis*. *Perognathus parvus* composed 84.2% of the total catch, with *P. maniculatus*, *O. leucogaster*, and *R. megalotis* comprising 9.4, 3.4, and 2.9 percent, respectively (Table 1).

The low trapping success in the fall reflects reduced surface activity and the post-breeding mortality of *P. parvus*, the most abundant species. Summer peaks reflect the termination of reproduction in this species and the increased foraging of weaned young. With the exception of *R. megalotis*, species composition was comparable to small mammal populations inhabiting slightly higher elevations on the reservation, where ground cover consists of more native vegetation (O'Farrell et al. 1975). The greater percent of captures of harvest mice on my study area may reflect this species propensity for habitats with a mixture of native and introduced vegetation (Black and Frischknecht 1971). *Peromyscus maniculatus* was the only species taken throughout the year, although it showed considerable seasonal variation in the number of individuals trapped. *Perognathus*

parvus was conspicuously absent in the coldest winter months (December and January) and *R. megalotis* was not trapped in the fall months of September and October. *Onychomys leucogaster* was captured each month except February, but trapping success for this species and *R. megalotis* never exceeded 1 percent. Other rodent species, including the sagebrush vole, mountain vole, Townsend's ground squirrel, pocket gopher, and bushy-tailed wood rat occur on parts of the reservation but were absent on my study area.

Because of the unpredictability of precipitation and extremes in temperature, desert rodent populations can demonstrate considerable annual fluctuations. The pocket mouse has specifically adapted to this environment (Schreiber 1978a), and the other species, because of their eurytopic habits, are able to survive at low population levels. Even though total population numbers may exhibit large annual oscillations, the proportional distribution of species probably remains stable over the long term.

Energy Budgets and Ingestion Rates

Energy flow in the individual is a function of the temperature gradient between body temperature (T_B) and ambient temperature (T_A). Heat is lost from the body when $T_A < T_B$ and gained by the body when $T_A > T_B$. The rate of metabolism is inversely proportional to the temperature gradient at temperatures below thermoneutrality and directly proportional to temperatures above it. Small rodents, with a relatively large body surface to body weight ratio, gain heat from the environment and dissipation of this heat load against a thermal gradient would require evaporative cooling and subsequent water loss, a luxury desert rodents cannot afford. However, these nocturnal animals rarely encounter ambient temperatures that exceed body temperatures (Table 2), so energy expenditure is mainly from thermogenesis and activity. Females experience additional demands during pregnancy and lactation. Seasonal changes in the insulatory

TABLE 1. Monthly trapping results for the Hanford Study Area, Benton County, Washington.

Month	Trap days	Species							
		P. parvus		P. maniculatus		O. leucogaster		R. megalotis	
		No.	Trap success (%)	No.	Trap success (%)	No.	Trap success (%)	No.	Trap success (%)
1969									
Sep.	200	9	(4.5)	1	(<1)	0		0	
Nov.	400	17	(4.2)	1	(<1)	2	(<1)	1	(<1)
1970									
Mar. °	280	50	(17.8)	6	(2.0)	0		1	(<1)
Apr.	297	50	(16.8)	13	(4.4)	2	(<1)	0	
May	298	92	(30.9)	10	(3.3)	3	(1.0)	1	(<1)
June	2216	544	(24.5)	39	(1.7)	19	(<1)	8	(<1)
July	550	166	(33.2)	7	(1.4)	3	(<1)	4	(<1)
Aug.	450	40	(8.9)	8	(1.8)	4	(<1)	1	(<1)
Sep.	1050	48	(4.6)	10	(1.0)	5	(<1)	0	
Oct.	1648	34	(2.1)	8	(<1)	6	(<1)	0	
Nov.	1300	2	(<1)	11	(<1)	1	(<1)	3	(<1)
Dec.	600	0		3	(<1)	1	(<1)	3	(<1)
1971									
Jan.	750	0		7	(<1)	3	(<1)	3	(<1)
Feb.	900	3	(<1)	6	(<1)	0		9	(1.0)
Mar.	750	39	(5.2)	3	(<1)	1	(<1)	4	(<1)
Apr.	950	60	(6.3)	4	(<1)	0		3	(<1)
May	750	47	(6.3)	2	(<1)	0		2	(<1)
Aug.	900	37	(4.1)	0		0		0	
Total	14,289	1,238	(8.7)	139	(1.0)	50	(<1)	43	(<1)

*Heavy rain and strong winds recorded at trapping site

properties of the pelage influence metabolic rates, but in small mammals this effect is minimal. Therefore annual energy expenditures of individuals are primarily the result of reproduction.

Reithrodontomys megalotis: This rodent was the smallest of the four species captured and it had a scattered distribution on the study area. Although never abundant, it is an opportunist, which enables it to exploit a variety of microhabitats.

Pearson (1960) calculated resting metabolism ($\text{ml O}_2 \text{ g}^{-1} \text{ hr}^{-1}$) in this small cricetid as—

$$E_r = 11.41 - 0.27 T_b \quad (T_b < 24.5 \text{ C}) \quad (4)$$

Adding the increment for activity (2.9 ml $\text{O}_2/\text{g/hr}$, Chew and Chew, 1970) to E_r ,

$$E_a = 14.31 - 0.27 T_A \quad (5)$$

Harvest mice construct elaborate, well-insulated nests which reduce energy expended for thermoregulation at lower temperatures. Thus,

$$E_n = 9.2 - 0.18 T_b \quad (T_b < 24.5 \text{ C}) \quad (6)$$

Harvest mice may be gregarious during the colder months; if so, their metabolic costs would be effectively reduced. Without nesting material, huddling can reduce metabolic rates 27–39 percent (Pearson 1960, Trojan and Wojciechowska 1968). With a nest, huddling reduces energy expended in heat production by about 13 percent (Grodzinski and Gorecki 1967) and significantly lowers

food consumption (Gebczynska and Gebczynski 1971). I used the latter figure (13 percent) to determine the savings from huddling in a nest.

Although *R. megalotis* is active throughout the year, its surface activity is presumably reduced during the colder months to minimize thermal stress. A male, with a radioactive tag, was monitored for three nights in November (Table 3). The average time spent above ground was 3.3 hours. I accepted this time as representative of both fall and winter activity. This estimate is probably a maximum because individuals may go several days without any surface activity during inclement weather. In fact, both Pearson (1960) and Gaertner (1968) allude to hypometabolism and possible torpor in *Reithrodontomys*. In the spring and summer, food availability and a more energetically favorable microclimate probably extend surface activity. Activity during these seasons was estimated as 4 hours/night, a value also used by Pearson (1960).

Estimated annual energy expenditure for an individual harvest mouse is shown in Table 4. Daily cost for maintenance in males (mean weight 10.76 g) and females (mean weight 9.48 g) is 9.11 and 8.02 kcal, respectively. This amounts to 0.85 kcal/g/day, of which thermoregulation accounts for about 74 percent and activity for 26 percent. Maintenance costs were 25 percent lower in

TABLE 2. Microenvironmental temperatures at the Hanford Study Area.

Time interval	Length (days)	Midpoint	Mean temperature, °C		
			In burrow (T_b)	At soil surface	
				Day (T_d)	Night (T_n)
Jun 17–Jun 28	12	Jun 22	24.2	43.9	22.4
Jun 29–Aug 3	36	Jul 16	26.9	40.1	22.5
Aug 4–Sep 10	38	Aug 22	26.0	37.0	18.5
Sep 11–Oct 12	32	Sep 26	19.2	21.6	8.2
Oct 13–Nov 15	34	Oct 29	12.6	8.6	-1.2
Nov 16–Dec 5	20	Nov 25	10.6 ^a	-1.8 ^b	-5.1 ^b
Dec 6–Jan 1	27	Dec 19	8.3 ^a	-4.5	-7.3
Jan 2–Feb 4	34	Jan 18	5.6 ^a	0.3	-4.5
Feb 5–Mar 12	36	Feb 22	5.6 ^a	5.2	-3.6
Mar 13–Apr 11	30	Mar 27	10.0	18.3	2.4
Apr 12–May 11	30	Apr 26	16.5	29.8	11.0
May 12–Jun 16	36	May 29	19.7	32.2	15.7

^aTemperatures taken at 9 dm (Stone, et al., 1972)

^bBased on 14 days data

TABLE 3. Activity patterns and average amount of time spent on the surface by a male *Reithrodontomys megalotis*.

Date	Time out	Time in	Elapsed time	Total time ^o	
November 27-28	21:26	23:29	2h:03m	2h:03m	
	1:30	1:50	0h:20m	0h:20m	
	2:43	2:54	0h:11m	0h:11m	
	3:54	4:03	0h:09m	0h:09m	
	4:45	5:20	0h:35m	0h:35m	
	7:08	7:33	0h:25m	0h:18m	
			Total	3h:36m	
November 28-29	21:16	22:35	1h:19m	1h:01m	
	23:40	23:56	0h:16m	0h:16m	
	1:50	2:10	0h:20m	0h:20m	
	3:37	3:58	0h:21m	0h:21m	
	4:42	4:52	0h:10m	0h:10m	
	5:47	6:12	0h:25m	0h:23m	
			Total	2h:31m	
November 29-30	20:10	21:07	0h:57m	0h:57m	
	22:17	22:42	0h:25m	0h:25m	
	0:24	0:50	0h:26m	0h:26m	
	3:12	4:01	0h:49m	0h:49m	
	5:32	6:15	0h:43m	0h:43m	
	6:47	7:07	0h:20m	0h:20m	
			Total	3h:40m	
				Mean time . . .	3h:16m

^oElapsed time minus time for periodic retreats to the burrow

TABLE 4. Annual maintenance energy expenditure for *Reithrodontomys megalotis* at the Hanford Study Area. Body weight: males = 10.76 g, females 9.48 g.

Dates	Time interval	Days	Hours daily		Metabolic costs (kcal) ^a					
			At rest	Active	Male			Female		
					E _r	E _a	E _m	E _r	E _a	E _m
6/17-6/28		12	20	4	62.0	20.6	82.6	54.6	18.1	72.7
6/29-8/3		36	20	4	159.9	62.5	222.4	140.9	55.0	195.9
8/4-9/10		38	20	4	172.7	73.8	246.5	152.2	65.0	217.2
9/11-10/12		32	20.7	3.3	195.0	66.5	261.5	171.1	58.6	229.7
10/13-11/15		34	20.7	3.3	254.4	84.0	338.4	224.2	74.0	298.2
11/16-12/5		20	20.7	3.3	156.1	52.8	208.9	137.5	46.6	184.1
12/6-1/1		27	20.7	3.3	219.4	74.9	294.3	193.3	66.0	259.3
1/2-2/4		34	20.7	3.3	301.7	89.2	390.9	265.8	78.6	344.4
2/5-3/12		36	20.7	3.3	319.4	92.6	412.0	281.4	81.6	363.0
3/13-4/11		30	20	4	229.3	84.3	313.6	202.0	74.3	276.3
4/12-5/11		30	20	4	195.2	70.6	265.8	172.0	62.2	234.2
5/12-6/16		36	20	4	212.0	75.1	287.1	186.7	66.2	252.9
Annual total (Kcal/yr) without huddling					2477.1	846.9	3324.0	2181.7	746.2	2927.9
					(74.5)	(25.5)		(74.5)	(25.5)	
with huddling					2289.1	846.7	3135.8	2016.2	746.2	2762.4
					(73.0)	(27.0)		(73.0)	(27.0)	

^aBased on microenvironmental temperatures (Table 2)

the spring and summer than in the fall and winter. For a 9 g individual of this species in central California, Pearson (1960) estimated daily energy costs averaged 7.6 kcal. His lower value reflects both the smaller average weight and the higher microenvironmental temperatures encountered by that population.

The coefficient of digestibility of this species, based on the ash tracer method, was 0.856. After adding the energy cost of pre-adult growth (E_{g_3}), ingestion rate for a male was calculated as:

$$I_m = 3473/0.856 = 4057 \text{ kcal/yr} \quad (7)$$

Based on a diet of 5.92 kcal/g (mean caloric value of stomach material, Schreiber 1979), a male would consume 686 g per year or about 1.9 g per day.

Weight gains were determined for each growth stage. At birth harvest mice weigh 1.5 g (Svihla 1931). This was increased 0.4 g to account for embryonic tissues. Weaning weight, prorated from that for deer mice, was calculated as 5.4 g. Mean body weight for adult males and nongravid (NG) females was 10.12 g ($N=34$). Caloric density for adult tissue was 1.58 kcal/g (Schreiber and Johnson 1975); other caloric values and growth efficiencies were given in the methods. Mean litter size (n) was 3.6 ($N=8$), with females producing 1.11 litters per year (L). The annual ingestion rate for a female was calculated as:

$$I_f = (2928 + 318)/0.856 = 3791 \text{ kcal/yr} \quad (8)$$

These mice expend 16.7 percent of this growth energy between conception and birth, 35.7 percent between birth and weaning, and 47.6 percent after weaning. Females would annually consume 642 g or 1.8 g per day based on the above diet. Huddling during the colder months would reduce total energy expenditure 5.7 percent.

Peromyscus maniculatus: Deer mice were the second most abundant rodent in the study area (Table 1). Because this species is also active periodically throughout the year, their activity patterns are presumably similar to those determined for harvest mice. For E_r , I adjusted the minimum rate measured by McNab and Morrison (1963, Table 1 and Fig. 3) by 24.5 percent, as suggested by Chew and

Chew (1970), to obtain the average resting metabolism:

$$E_r = 9.3 - 0.2 T_b \quad (T_b > 27.1 \text{ C}) \quad (9)$$

Correcting this for the energy used during activity:

$$E_a = 12.2 - 0.2 T_A \quad (10)$$

The insulating effects of a nest reduce E_r to:

$$E_n = 7.4 - 0.13 T_b \quad (T_b < 27.1 \text{ C}) \quad (11)$$

Daily, males require about 14.1 kcal and females 12.1 kcal for maintenance (Table 5) or 0.63 kcal/g/day in the spring and summer.

Adding the growth increment and using a coefficient of digestibility of 0.879 calculated for animals living in the wild, the ingestion rate for males was:

$$I_m = 5344/0.879 = 6080 \text{ kcal/yr} \quad (12)$$

In addition to maintenance costs, females in the study areas produced an average of 1.32 litters per year with a mean number of 4.7 young ($N=44$). At birth, deer mice weigh an average of 1.8 g (Svihla 1934); embryonic tissues add 1.1 g. Weaning takes place in about 25 days, when animals weigh about 11 g (Svihla 1934, Chew and Chew 1970). Mean adult body weight of males and NG females was 17.5 g ($N=162$) and adult tissue of this species has a caloric value of 1.56 kcal/g (Schreiber and Johnson 1975). Previously cited values were used for other tissue energy and growth efficiencies. Ingestion rate for females was:

$$I_f = (4425 + 753)/0.879 = 5891 \text{ kcal/yr} \quad (13)$$

Of the 753 kcal used for growth, 16.9 percent was used before birth, 56.1 percent from birth to weaning, and 26.9 percent after weaning. Deer mice on the study area consumed diets with a mean caloric value of 5.75 kcal/g (Schreiber 1979). Therefore, to meet the required energy demands, males would consume 1.06 kg/yr and females 1.03 kg/yr, or about 2.9 g/day. This amounts to 14.6 percent and 16 percent of the body weight in males and females, respectively. Johnson and Groepper (1970) estimated a 20 g deer mouse in the North Plains consumed 1.9 g of food daily or about 9 percent of its body weight. Deer mice on standard rations at temper-

atures 10–15 C reportedly consumed an average of 3 g/day (Hatfield, 1940) or 2.8 g/day at 28 C (Sealander 1952).

Energy requirements effecting reproductive stress may substantially increase (e.g., 200 percent) between birth and the end of weaning (Stebbins 1977), so my estimates for females may be somewhat low. Besides huddling, which effectively reduces metabolic costs 5.5 percent (Table 5), deer mice no doubt also resort to additional means of reducing energy expenditure during periods of stress. Marten (1973) found this species may sharply reduce its activity through the summer and activity becomes compressed into the early part of the night, when conditions probably permit a more favorable heat exchange. For the same reason, diurnality may increase in colder months. In winter, pelage (insulation) increases and also contributes to a lower metabolism. Howard (1951) suggested huddling is an important part of energy reduction in deer mice, with torpor occurring in grouped animals. Torpor in this species has also been observed by others (e.g., Morhardt and Hudson 1966, Morhardt 1970). Additionally, Kritzman (1974) and O'Farrell (1975a) suggested possible summer estivation or hypothermia for animals at the Hanford Reservation. Although the rhythmicity of torpor in natural populations is unknown, it could

serve both to conserve water in the summer and reduce energy costs at low temperatures or during periods of food scarcity. Hart (1958) also suggested possible alterations in the ability to metabolize food during such periods of stress. As an additional factor, food caches may offer a significant buffer during periods of severe weather and with sufficient stored food, individuals would spend less time on the surface exposed to unfavorable micro-environmental temperatures.

Peromyscus maniculatus is omnivorous, demonstrating marked seasonal variation in its diet (Johnson 1964). On other parts of the reservations this species relied heavily on insects for food from spring until fall (Kritzman 1974), probably switching to a more granivorous diet as this food source diminished. It would appear then that competition for available seeds between this species and the more abundant *P. parvus* would be reduced due to their periods of activity, deer mice becoming more dependent on seeds in the colder months when pocket mice are dormant. Additional research is necessary to properly evaluate these aspects of deer mice energetics.

Onychomys leucogaster: At an average weight of 25 g, the stout-bodied northern grasshopper mouse was the largest rodent species on the study area. It was present in

TABLE 5. Annual maintenance energy expenditure for *Peromyscus maniculatus* at the Hanford Study Area. Body weight: males = 19.17 g, females 16.83 g.

Dates	Time interval	Hours daily		Metabolic costs (kcal) ^a					
		At rest	Active	Male			Female		
				E _r	E _a	E _m	E _r	E _a	E _m
6/17-6/28	12	20	4	95.0	34.2	129.2	83.4	30.0	113.4
6/29-8/3	36	20	4	258.4	102.0	360.4	226.8	89.6	316.4
8/4-9/10	38	20	4	283.2	118.9	402.1	248.6	104.4	353.0
9/11-10/12	32	20.7	3.3	301.7	102.5	504.2	264.9	90.0	354.9
10/13-11/15	34	20.7	3.3	378.2	128.5	506.7	332.0	112.8	444.8
11/16-12/5	20	20.7	3.3	232.4	80.3	312.7	204.0	70.5	274.5
12/6-1/1	27	20.7	3.3	329.1	111.9	441.0	289.0	98.2	387.2
1/2-2/4	34	20.7	3.3	433.9	135.2	569.1	380.9	118.7	499.6
2/5-3/12	36	20.7	3.3	459.4	141.0	600.4	403.3	123.8	527.1
3/13-4/11	30	20	4	336.8	130.0	466.8	295.7	114.1	409.8
4/12-5/11	30	20	4	292.6	110.4	403.0	256.9	96.9	353.8
5/12-6/16	36	20	4	324.6	120.6	445.2	285.0	105.8	390.8
Annual total (Kcal/yr) without huddling				3825.3	1315.5	5140.8	3270.5	1154.8	4425.3
				(74.4)	(25.6)		(73.9)	(26.1)	
with huddling				3547.8	1315.5	4863.3	3026.9	1154.8	4181.7
				(73.0)	(27.0)		(72.4)	(27.6)	

^aBased on microenvironmental temperatures (Table 2)

low numbers throughout the year. Traps holding these mice were often adjacent to those with mutilated pocket mice, indicating *O. leucogaster's* carnivorous tendency and predatory feeding habits. To my knowledge, no metabolic-temperature function equation presently exists for this species, so I used the equation for *O. torridus* (Chew and Chew 1970), a species of similar size. Average resting metabolism is:

$$E_r = 7.24 - 0.17 T_b \quad (T_b > 27.1 \text{ C}) \quad (14)$$

This species uses a nest (Ruffer 1965) and, assuming Pearson's (1960) correction for its insulating properties,

$$E_n = 5.86 - 2.12 T_b \quad (T_b < 27.1 \text{ C}) \quad (15)$$

above-ground activity increases energy expenditure; so,

$$E_a = 10.14 - 0.17 T_A \quad (16)$$

presumably, this species has periods of activity similar to harvest mice and deer mice. Annual maintenance costs were estimated as 4857 and 5215 kcal for males and females, respectively (Table 6). Generally, metabolic costs are about 25 percent greater in the colder months than in the warmer months. Nest burrows are shared by male-female pairs (Ruffer 1965); such huddling would produce a 5 percent savings in energy.

Males digest an average of 90.3 percent of their caloric diet. $E_{g_3} = 303$ kcal for this species, so ingestion rate was calculated as:

$$I_m = 5160/0.903 = 5714 \text{ kcal/yr} \\ \text{(without huddling)}$$

or

$$= 4893/0.903 = 5419 \text{ kcal/yr} \\ \text{(with huddling)} \quad (17)$$

Grasshopper mice weigh 2.2 g at birth (Svihla 1936), excluding embryonic tissues (~ 0.6 g). Females raised an average of 1.07 litters per year with a mean litter of 3.3 young ($N=14$). Young are weaned at about 23 days at an average weight of 13.2 g (Pinter 1970) and the mean weight of mature males and NG females was 22.6 g. Using a caloric value of 1.61 kcal/g for adult tissue (Schreiber and Johnson 1975) and other values cited earlier, energy cost for growth from conception to subadult is 660 kcal. The majority of this growth energy is expended after birth. Prenatal growth accounted for 10.6 percent, weaning period for 43.5 percent, and postweaning growth for 45.9 percent. With a digestibility coefficient of 0.892, ingestion rate for females is:

$$I_f = (5215 + 660)/0.892 = 6587 \text{ kcal/yr} \\ \text{(without huddling)}$$

or

TABLE 6. Annual maintenance energy expenditure for *Onychomys leucogaster* at the Hanford Study Area. Body weight: males = 24.3 g, females 26.2 g.

Time interval		Hours daily		Metabolic costs (Kcal) ^a					
Dates	Days	At rest	Active	Male			Female		
				E_r	E_a	E_m	E_r	E_a	E_m
6/17-6/28	12	20	4	82.6	35.6	118.2	89.0	38.3	127.3
6/29-8/3	36	20	4	219.2	105.0	324.2	236.3	113.2	349.5
8/4-9/10	38	20	4	239.3	124.1	363.4	258.0	133.8	371.2
9/11-10/12	32	20.7	3.3	274.3	107.2	381.5	295.7	115.5	411.2
10/13-11/15	34	20.7	3.3	357.1	134.8	491.9	385.0	145.3	530.3
11/16-12/5	20	20.7	3.3	222.1	83.9	306.0	239.5	90.5	330.0
12/6-1/1	27	20.7	3.3	316.2	117.4	433.6	340.9	126.6	467.5
1/2-2/4	34	20.7	3.3	426.9	142.6	569.5	460.2	153.8	614.0
2/5-3/12	36	20.7	3.3	452.0	147.6	599.6	487.3	159.1	646.4
3/13-4/11	30	20	4	325.4	135.1	460.5	350.9	145.6	495.5
4/12-5/11	30	20	4	272.9	115.5	388.4	294.3	124.5	418.8
5/12-6/16	36	20	4	293.9	126.0	419.9	316.9	135.8	452.7
Annual total (Kcal/yr) without huddling				3481.9	1374.8	4856.7	3733.7	1482.0	5215.4
				(71.7)	(28.3)		(71.6)	(28.4)	
with huddling				3215.6	1374.8	4590.4	3446.3	1482.0	4928.3
				(70.1)	(29.9)		(69.9)	(30.1)	

^aBased on microenvironmental temperatures (Table 2)

$$(4928 + 660)/0.892 = 6265 \text{ kcal/yr} \\ \text{(with huddling)} \quad (18)$$

Males and females which huddle must daily digest 15.6 (0.69) and 17.2 kcal (0.76 kcal/g), respectively. Diets of grasshopper mice include a variety of insects and seeds plus some animal flesh and green vegetation. Stomach material of this species at Hanford had a mean caloric value of 5.22 kcal/g (Schreiber 1979), about 8 percent lower than the value reported for North Plains individuals (Johnson and Groepper 1970). Based on this caloric diet each male and female in the population studied would annually consume about 1038 g (2.9) and 1200 g (3.3 g/day) of food, respectively. By comparison, in the laboratory on diets of beef liver (72 percent water), grasshopper mice consumed 4.1 g/day (Whitford and Conley 1971).

The incidence of seeds in the diet of this species may significantly increase in the fall and winter (Flake 1971), suggesting it relies on seed caches when insects are less available. Less surface activity would decrease metabolic demands in the colder months. To facilitate this energy savings individuals may restrict more of their winter "hunting" activity to burrow systems. Burrows serve as hibernacula for insects, lizards, and torpid mice, all potential food sources for *O. leucogaster*. The generally high fat deposits of this

rodent probably conserve body heat when it is forced to be active on the surface during the cold months.

Ecological Relationships

Individually, *P. maniculatus* and *O. leucogaster* process between 2½ to 3 times as much energy during the year as *P. parvus*, the most common species (Table 7). The smaller size of the pocket mouse and its periodic use of torpor account for this difference. Based on the relative numbers of individuals (Table 1), however, *P. parvus* dominates as the primary "energy mover" in the small mammal community of this cold desert ecosystem. At the population level, the annual contribution of pocket mice to community energy exchange is nearly 4 times that of deer mice, 11 times that of grasshopper mice, and about 17 times that of harvest mice. Even at high population levels, however, the granivorous pocket mouse does not significantly affect its primary food resource, cheatgrass (Schreiber 1978b). Thus, it is reasonable to assume the euryphagic and less abundant species represented in this paper would also have an insignificant impact on their diverse food resources.

These results additionally refine our understanding of energetics of small mammal com-

TABLE 7. Estimated annual ingestion rates of selected Great Basin rodents.

Species	Sex	E_m^a (kcal/yr)	Coefficient of digesti- bility	Mean litter size	Number litters per year	Energy for growth (kcal)	Ingestion rate	
							(kcal/yr)	(kg/yr) ^b
<i>Perognathus</i> <i>parvus</i>	M	2010	0.892			205	2483	0.50
	F	1774	0.911	3.9	1.10	360	2342	0.47
<i>Peromyscus</i> <i>maniculatus</i>	M	5141	0.879			203	6080	1.06
	F	4425	0.879	4.7	1.32	753	5891	1.02
<i>Onychomys</i> <i>leucogaster</i>	M	4857	0.903			303	5714	1.09
	F	5215	0.892	3.3	1.07	660	6587	1.26
<i>Reithrodontomys</i> <i>megalotis</i>	M	3324	0.856			148	4057	0.68
	F	2928	0.856	3.6	1.11	318	3791	0.64

^aMaintenance energy (E_m) is calculated without the advantage of savings from huddling.

^bBased on the mean caloric values of diets of wild mice (Schreiber, 1979)

^cTaken from Schreiber (1978b)

munities. Unfortunately, the question of the ecological significance of these consumers remains unanswered and a matter of considerable discussion (e.g., Naumov 1975, Chew 1978). As deserts come under increasing pressure for development, the understanding of community function and of the relative cost/benefit of perturbing its various components becomes imperative. Research must now focus on integrating information on bioenergetics with other parameters, such as resource allocation, nutrient cycling and inter- and intraspecific competition, and developing new hypotheses on the role of small mammal consumers.

ACKNOWLEDGMENTS

I am indebted to Dr. Donald R. Johnson, University of Idaho, who initiated this project and secured its financial support. Additional support and facilities were provided by the Joint Center for Graduate Studies and Battelle Pacific Northwest Laboratories, Richland, Washington (USAEC Contract AT(45-1)-2042). The research represents a portion of my dissertation, submitted in partial fulfillment of the requirements for a Doctor of Philosophy degree in zoology from the University of Idaho. Final typing was provided by Sally Vreeland.

LITERATURE CITED

- BLACK, H. L., AND N. C. FRISCHKNECHT. 1971. Relative abundance of mice on seeded sagebrush-grass range in relation to grazing. U.S.D.A. Forest Serv. Res. Note INT-147. 8 p.
- CHEW, R. M. 1978. The impact of small mammals on ecosystem structure and function, p. 167-180. In: D. P. Snyder (ed.) Populations of small mammals under natural conditions. Spec. Publ. Series, Vol. 5, Pymatuning Laboratory of Ecology, Univ. Pittsburgh, Linesville, Pennsylvania. 237 p.
- CHEW, R. M., AND A. E. CHEW. 1970. Energy relationships of the mammals of a desert shrub (*Larrea tridentata*) community. Ecol. Monogr. 40:1-21.
- DAUBENMIRE, R. 1970. Steppe vegetation of Washington. Washington State Agri. Exp. Sta. Tech. Bull. 62, Coll. of Agri., Washington State Univ., 131 p.
- DROZDZ, A. A., GORECKI, AND K. SAWICKA-KAPUSTA. 1972. Bioenergetics of growth in common voles. Acta Theriol. 17:245-257.
- FLAKE, L. 1971. An ecological study of rodents in a short-grass prairie of northeastern Colorado. U.S./IBP, Grasslands Biome Tech. Rep. 100. Colorado State Univ., Fort Collins. 118 p.
- FRENCH, N. R., B. G. MAZA, AND A. P. ASCHWANDERN. 1966. Periodicity of desert rodent activity. Science 154:1194-1195.
- GAERTNER, R. A. 1968. Seasonal variations in the energy budgets of the harvest mouse, *Reithrodontomys fulvescens*, and the cotton rat, *Sigmodon hispidus*. Unpublished dissertation. Univ. of Arkansas. 149 p.
- GEBCZYNSKA, Z., AND M. GEBCZYNSKI. 1971. Insulating properties of the nest and social temperature regulation in *Clethronomys glareolus* (Schreber). Ann Zool. Fennici 8:104-108.
- GORECKI, A. 1965. Energy values of body in small mammals. Acta Theriol. 10:333-352.
- GRODZINSKI, W., AND A. GORECKI. 1967. Daily energy budgets of small rodents, p. 295-314. In: K. Petruszewicz, (ed.), Secondary productivity of terrestrial ecosystems. Panstwowe Wydawnictwo Naukowe, Poland.
- HANSSON, L. 1967. Index line catches as a basis of population studies on small mammals. Oikos 18:261-176.
- HART, J. S. 1958. Metabolic alterations during chronic exposure to cold. Fed. Proc. 17(4):1045-1054.
- HATSFIELD, D. M. 1940. Activity and food consumption in *Microtus* and *Peromyscus*. J. Mammal. 21:29-36.
- HEDLUND, J. D., D. T. McCULLUGH, AND W. H. RICKARD. 1975. Mouse populations on Knob and Kettle topography in south-central Washington. Northwest Sci. 49:253-260.
- HOWARD, W. E. 1951. Relation between low temperature and available food to survival of small mammals. J. Mammal. 32:300-312.
- JOHNSON, D. R. 1964. Effects of range treatment with 2, 4-D on food habits of rodents. Ecol. 45:241-249.
- JOHNSON, D. R., AND K. L. GROEPPER. 1970. Bioenergetics of North Plains rodents. Amer. Midland Nat. 84:537-548.
- KACZMARSKI, F. 1966. Bioenergetics of pregnancy and lactation in the bank vole. Acta Theriol. 14:167-179.
- KRITZMAN, E. B. 1974. Ecological relationships of *Peromyscus maniculatus* and *Perognathus parvus* in eastern Washington. J. Mammal. 55:172-188.
- McNAB, B. K. 1963. A model of the energy budget of a wild mouse. Ecol. 44:521-532.
- McNAB, B. K., AND P. R. MORRISON. 1963. Body temperature and metabolism in subspecies of *Peromyscus* from arid and mesic environments. Ecol. Monogr. 33:63-82.
- MARTEN, G. G. 1973. Time patterns of *Peromyscus* activity and their correlations with weather. J. Mammal. 54:169-188.
- MIGULA, P. 1969. Bioenergetics of pregnancy and lactation in European common vole. Acta Theriol. 14:167-179.
- MORHARDT, J. E. 1970. Body temperatures of white-footed mice (*Peromyscus* sp.) during daily torpor. Comp. Biochem. Physiol. 33:423-439.
- MORHARDT, J. E., AND J. W. HUDSON. 1966. Daily torpor induced in white-footed mice (*Peromyscus* sp.) by starvation. Nature 212:1046-1047.

- MURIE, M. 1961. Metabolic characteristics of mountain, desert, and coastal populations of *Peromyscus*. *Ecol.* 42:723-740.
- MYRCHA, A., AND W. WALKOWA. 1968. Changes of the caloric value of the body during the postnatal growth development of white mice. *Acta Theriol.* 13:391-400.
- NAUMOV, N. P. 1975. The role of rodents in ecosystems of the northern deserts of Eurasia, p. 299-309. In: F. B. Golley, et al. (eds.), *Small mammals: their productivity and population dynamics*. Intern. Biol. Prog. 5, Cambridge Univ. Press, Cambridge. 451 p.
- O'FARRELL, T. P. 1975a. Small mammals, their parasites and pathologic lesions on the Arid Lands Ecology Reserve, Benton County, Washington. *Amer. Midl. Nat.* 93:377-387.
- . 1975b. Seasonal and altitudinal variations in populations of small mammals on Rattlesnake Mountain, Washington. *Amer. Midl. Nat.* 94:190-204.
- O'FARRELL, T. P., R. J. OLSON, R. O. GILBERT, AND J. D. HEDLUND. 1975. A population of Great Basin pocket mice, *Perognathus parvus*, in the shrub-steppe of south-central Washington. *Ecol. Monogr.* 45:1-28.
- PEARSON, O. P. 1960. The oxygen consumption and bioenergetics of harvest mice. *Physiol. Zool.* 33:152-160.
- PETTICREW, B. G., AND A. M. F. S. SADLER. 1970. The use of index trap lines to estimate population numbers of deer mice (*Peromyscus maniculatus*) in a forest environment in British Columbia. *Canadian J. Zool.* 48:385-389.
- PINTER, A. J. 1970. Reproduction and growth for two species of grasshopper mice (*Onychomys*) in the laboratory. *J. Mammal.* 51:236-243.
- RUFFER, D. G. 1965. Burrows and burrowing behavior of *Onychomys leucogaster*. *J. Mammal.* 46:241-247.
- SCHREIBER, R. K. 1973. Bioenergetics of rodents in the northern Great Basin desert. Unpublished dissertation, Univ. Idaho, Moscow. 133 p.
- . 1978a. Paleoecology of desert vegetation and rodents in the Great Basin. *Biologist* 60:19-33.
- . 1978b. Bioenergetics of the Great Basin pocket mouse, *Perognathus parvus*. *Acta Theriol.* 23:469-488.
- . 1979. Coefficients of digestibility and caloric diet of rodents in the northern Great Basin desert. *J. Mammal.* 60: (In press).
- SCHREIBER, R. K., AND D. R. JOHNSON. 1975. Seasonal changes in body composition and caloric content of Great Basin rodents. *Acta Theriol.* 20:343-364.
- SEALANDER, J. A., JR. 1952. Food consumption in *Peromyscus* in relation to air temperature and previous thermal experience. *J. Mammal.* 33:206-218.
- SOHOLT, L. F. 1973. Consumption of primary productivity by a population of kangaroo rats (*Dipodomys merriami*) in the Mojave Desert. *Ecol. Monogr.* 43:357-376.
- STEBBINS, L. L. 1977. Energy requirements during reproduction of *Peromyscus maniculatus*. *Can. J. zool.* 55:1701-1704.
- STONE, W. A., D. E. JENNE, AND J. M. THORP. 1972. Climatology of the Hanford Area. BNWL-1605, June 1972. Battelle Pacific Northwest Lab., Richland, Washington.
- SVIHLA, A. 1934. Development and growth of deer mice (*Peromyscus maniculatus artemisiae*). *J. Mammal.* 15:99-104.
- SVIHLA, R. D. 1931. Notes on desert and dusky harvest mice (*Reithrodontomys megalotis megalotis* and *Reithrodontomys megalotis nigrescens*). *J. Mammal.* 12:363-365.
- . 1936. Breeding and young of the grasshopper mouse (*Onychomys leucogaster fuscogriseus*). *J. Mammal.* 17:172-173.
- TROJAN, P., AND B. WOJCIECHOWSKA. 1968. The effects of huddling on resting metabolism rate in European common vole *Microtus arvalis*. *Bull. Acad. Pol. Sci. Cl. II.* 16:107-109.
- WHITFORD, W. G., AND M. I. CONLEY. 1971. Oxygen consumption and water metabolism in a carnivorous mouse. *Comp. Biochem. Physiol.* 40:797-803.
- WIENER, J. G., AND M. H. SMITH. 1972. Relative efficiencies of four small mammal traps. *J. Mammal.* 53:868-873.



Schreiber, R K. 1979. "ANNUAL ENERGY BUDGETS FOR 3 COMMON RODENT SPECIES IN THE NORTHERN GREAT BASIN WASHINGTON USA." *The Great Basin naturalist* 39, 143–154.

View This Item Online: <https://www.biodiversitylibrary.org/item/33879>

Permalink: <https://www.biodiversitylibrary.org/partpdf/91092>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Brigham Young University

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.