

REFERENCES

- Audas, J. W. (1950) *The Australian Bushland*. W. A. Hamer: N. Melbourne.
- Bentham, G. (1870) *Flora Australiensis*. Vol. 5, L. Reeve and Co.: London.
- Brown, R. (1814) *Botany of Terra Australia*. In M. Flinders. *Voyage to Terra Australia*. Vol. 2 Nicol: London.
- Brown, R. (1830) *supplementum primum prodromi florum Novae Hollandiae*. R. Taylor: London.
- Burke, I. and K. (eds.) (1967) *The Spanish at Port Jackson*. Australian Facsimile Society: Sydney.
- Cavanilles, A. J. (1800) Observations on the soil, natives and plants of Port Jackson and Botany Bay. *Anales de Hist. Nat.* 3. 181-245. *Trans. R. J. Dorr*.
- Flinders, M. (1814) *A Voyage to Terra Australis; undertaken . . . in the Year 1801, 1802 and 1803* 2 vols. Nicol: London.
- Mack, J. D. (1966) *Matthew Flinders 1774-1814* Nelson: Australia.
- Meissner, K. F. (1854) *Proteaceae. Linneaea* 26 : 352-3.
- Mueller, F. (1853) *First General Report. Votes and Proceedings of the Legislative Council 1853-4* Vol. 1 Part 2, p 897. Government Printer : Melbourne.
- Stearn, W. T. (1960) *Introduction to facsimile edition of R. Brown's Prodrum florum Novae hollandiae*. In J. Cramer and H. J. Swann (eds.) *Historiae naturalist classica* 6. Cramer : Weinheim.

Little Red Flying-foxes Collecting Water in Fur

BY RICHARD H. LOYN¹

During a period of hot weather in January 1979 a flock of about 200 Little Red Flying-foxes (*Pteropus scapulatus*) was found roosting in 14m high River Red Gum (*Eucalyptus camaldulensis*) regrowth near Barmah Lake on the River Murray. At dusk the bats left the roost and began flying low over the water on a tributary creek, gliding and making loud splashes as they belly-flopped onto the surface. The heads were held high and did not touch the surface. By watching individual bats I could see that after doing this each would fly to a nearby tree, hang upside down and lick the water vigorously from the fur on its chest. I concluded that the function of this behaviour was to obtain water for drinking rather than washing as the bats did not appear to lick parts which had not already been wetted, and their movements suggested that they were swallowing substantial volumes of water.

It is well known that bats of many

species fly low over water to obtain insects or water. However, I am not aware of any reports of collecting water in the belly fur. Troughton (1973) mentions that fruit-bats "drink regularly, usually while on the wing; lapping the water as they glide over its surface." Perhaps this is a mistaken interpretation of the belly-flopping method. However, I do not know if flying foxes use the method regularly or only on hot days. The temperature had reached 45°C on the day of observation (5.1.79) and the bats may have been under stress (and in need of extra water); three dead ones were found under the roost. The most obvious food source was the blossoms of River Red Gums which were flowering nearby.

Acknowledgement

I am grateful to Miss Joan Dixon and staff of the National Museum, Victoria for confirming identity of one of the dead bats.

REFERENCE

- Troughton, E. 1973 *Furred Animals of Australia*. 9th Edition (1967), revised 1973. Angus and Robertson, Sydney. 314 pp.

1. Mountain Forest Research Station, Sherbrooke, Vic., 3789.



Loyn, Richard H. 1981. "Little Red Flying-foxes Collecting Water in Fur." *The Victorian Naturalist* 98, 194–194.

View This Item Online: <https://www.biodiversitylibrary.org/item/123293>

Permalink: <https://www.biodiversitylibrary.org/partpdf/92427>

Holding Institution

Field Naturalists Club of Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Permissions to digitize granted by rights holder.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.