

THE GEOGRAPHICAL DISTRIBUTION OF SOUTHERN AFRICAN HYDROIDS

By

N. A. H. MILLARD

South African Museum, Cape Town

(With 9 figures, 8 tables and 2 appendices)

[MS. accepted 9 August 1977]

ABSTRACT

The geographical distribution of 251 species of hydroid Cnidaria round the coast of southern Africa is analysed by means of a system of radial sectors. It is concluded that there is no distinct west coast fauna as such, but that a boundary zone exists at approximately 31°S latitude between an east coast region with mainly tropical affinities and a temperate west-plus-south coast region with a reduced tropical component and a large endemic population. The main centre of the endemic population is the Agulhas Bank. The relationships of the tropical species are mainly with the Western Indian Ocean Province and the Indo-West-Pacific Region. Deep-water species (45) which occur below 400 m are of a mixed nature and show no clear relationship with other parts of the world. Deep-water species are classified into stenobathic/eurybathic and stenothermic/eurythermic categories, and their distribution relative to water temperatures is discussed.

CONTENTS

	PAGE
Introduction	159
Methods	160
Faunal boundaries	164
The tropical component	169
The temperate component	170
The endemic component	173
The cosmopolitan and scattered components	174
Distribution and depth	175
Estuarine and brack-water species	179
Conclusion	182
References	183
Appendix 1	185
Appendix 2	188

INTRODUCTION

This paper is an attempt to analyse the distribution of the hydroid fauna of the coast of southern Africa as documented in a recent monograph (Millard 1975). Since the latter went to press a few more taxonomic papers on, or lists of, hydroids have appeared, namely Arnaud *et al.* (1976), Beurois (1975), Calder (1975, 1976), Cooke (1975), Cornelius (1975a, 1975b), Mergner & Wedler (1977), Millard & Bouillon (1975), Millard (1977a) and Watson (1975). Among

these are records which extend the range of some South African species and they have been taken into account in the analysis which follows, as also have records from Buchanan (1957), Michel (1974), Blanco (1967, 1968, 1973, 1974) and Blanco & Bellusci de Miralles (1972a, 1972b), which have only recently become available to the author.

Within the borders of southern Africa, here taken to be the 20° parallel of south latitude, several new distribution records from deep water off the Natal coast and some new records for the country were published by Millard (1977b). There are also scattered identifications by the author, as yet unpublished, which are listed in Appendix 1.

The following changes in name have been adopted here:

Antennella africana Broch, 1914, to *A. quadriaurita* Ritchie, 1909, *fide* Millard (1977b: 123).

Eulaomedeia calceolifera (Hincks, 1871) to *Campanularia calceolifera* Hincks, 1871, *fide* Cornelius (1975a: 254).*

Lafoea fruticosa (M. Sars, 1851) to *Lafoea dumosa* (Fleming, 1820) *fide* Cornelius (1975b: 385).

Obelia bicuspidata Clarke, 1875, to *O. bidentata* Clarke, 1875, *fide* Cornelius (1975a: 260).

Any zoogeographical analysis is aimed at pinpointing population changes within an area, and, if possible, relating them to physical or chemical factors and determining relationships with other areas. Such an analysis is subject to certain unavoidable weaknesses—the assignment of species to categories or components is largely subjective; numbers of species within categories are liable to change as more records are forthcoming (for instance some of the South African endemics may in time be expected to turn up outside our boundaries); and numbers of species may be influenced by different intensities of collecting in different areas or by misidentifications. The longer we wait the more reliable will be the results, but results, if only interim ones, are needed now. The larger the number of species involved the more meaningful will be the conclusions. One must bear in mind that it is the main trends which are significant rather than the details.

METHODS

The first step in the analysis was to remove all doubtful records and incomplete identifications. This left a total of 251 species.

The first objective was an analysis of distribution within the borders of southern Africa, and for this some system of comparison was necessary. When dealing with the littoral zone it is a comparatively simple matter to convert the distance along the coast to a straight line, a method with obvious practical

* Cornelius pointed out that the type species of *Campanularia* is *Sertularia verticillata* Linnaeus, 1758, and not *Sertularia volubilis* Linnaeus, 1758, as previously thought (Millard 1975: 203). Since *S. verticillata* is a branched species the diagnosis of *Campanularia* given previously (Millard 1975: 203) must be altered to include branching forms, and *Eulaomedeia* sunk in *Campanularia*.




Fig. 1. Map of southern Africa showing the main collecting localities. From Millard (1975).

advantages, and this has been done by Stephenson (1947), Stephenson & Stephenson (1972), Day (1967) and others. However, when dealing with the sublittoral area as well, this is clearly not possible for a continent of the shape of southern Africa. After much consideration the following arbitrary method was devised, in which the coastal waters were divided into thirty-six sectors of approximately equal area (Fig. 2).

The central point for the sectors was chosen as the intersection of the latitude line of 20°S and the longitude line of 24°E , a point which is approximately mid-way between the west and east coastlines. From this central point radii were drawn at 5° intervals, starting with one through the Cape Peninsula, and then working outwards in both directions. The sectors thus constructed were numbered 1–36. The reason for placing one radius (Number 14/15) through the Cape Peninsula is that Cape Point is generally recognized as an approximate zoogeographical boundary between the cold Benguela water on the west and the warmer Agulhas water on the east, and might be expected to have some significance. It so happens that another radius (Number 15/16) conveniently passes through Cape Agulhas, the most southerly point of Africa, and a third (Number 23/24) through the boundary between Transkei and Natal.




Fig. 2. Map of southern Africa illustrating the sector system which was used to analyse the distribution.

It must be emphasized that this is an arbitrary method, and obviously could not be applied to all coastlines, but it is felt that it might form a useful working method and basis for comparison with other southern African marine groups. It has the advantage that each sector covers an approximately equal area of sea and 'straight-line' length of coast and represents a cross-section through all depths. Since the sectors are roughly normal to the coast and the currents tend to follow the coast, each sector is affected in a similar way and population differences related to water temperature should be highlighted. It is felt that the advantages of this method at least outweigh the disadvantages (and impracticabilities) of the 'straightening-out' method of Stephenson.

The second objective was an analysis of the world-wide distribution of the southern African species, and for this the species were delegated to components defined as follows:

TROPICAL

Circumtropical. Present in the tropics of all three oceans, but may spread north or south into warm temperate seas as well.

Indo-West-Pacific Region. Stretching from the Suez Canal across the Indian Ocean and the East Indies into the Western Pacific as far as Hawaii and Easter Island. Includes northern Australia, Madagascar and East Africa. May spread into warm temperate seas as well.

Western Indian Ocean Province. The western section of the Indo-West-Pacific Region, bounded in the north by the Persian Gulf, and including Madagascar and surrounding islands.

TEMPERATE. (In a trial analysis it was found impossible to differentiate between cold temperate and warm temperate fauna when assessing world distribution.)

Antitropical (bipolar). Present in the temperate zones of all three oceans, and present in both the Northern and the Southern hemisphere, but not in the tropics except perhaps just at the edge.

Southern. Restricted to the temperate waters of the Southern hemisphere, but often in the subantarctic islands as well, and rarely reaching the Antarctic.

Atlantic. Restricted to the northern and/or southern temperate zones of the Atlantic, but absent from the tropics. May spread further north into the subarctic.

ENDEMIC

OTHER

Cosmopolitan. Present in all three oceans, and occurring in both tropical and temperate seas.

Scattered. Present in various regions, but not sufficiently widespread to be included in Cosmopolitan. Does not fit any other category.

A list of species, together with their components and distribution within thirty-four sectors, is given in Appendix 2. (Sectors 35 and 36 have been omitted as no records were available from this region.)

FAUNAL BOUNDARIES

It has long been recognized that the marine fauna of southern Africa includes both tropical elements, predominating on the east coast, and temperate elements, predominating on the south and west coasts. Most authorities, following Stephenson (1947) and Stephenson & Stephenson (1972), have recognized a separate east coast (tropical or subtropical) fauna, a south coast fauna (warm temperate) and a west coast fauna (warm or cold temperate) for the littoral region. These earlier views are summarized by Briggs (1974), who includes the east coast in the Western Indian Ocean Province of the Indo-West-Pacific tropical region; the south coast in the Agulhas Province of the South African warm temperate region; and the west coast as a separate South West Africa Province of the South African warm temperate region. Opinions have differed as to the position of the boundary between the east and south coast, and whether the west coast represents a separate faunal province, and if so whether it should be regarded as cold temperate or warm temperate.

The distribution of the faunal components of the continental shelf and the extent of the faunal regions is largely dependent on the ocean currents round the South African coast, in general the warm southward-flowing Agulhas Current being responsible for the southerly extension of tropical forms on the east coast, and the cold northward-flowing Benguela Current being responsible for the more northerly distribution of cool-water forms on the west coast. It is not intended to discuss these current systems in any detail here since accounts are available in most textbooks of marine biology (a concise general account is given by Stephenson & Stephenson (1972) and more recent contributions are summarized by M. J. Penrith (1976)).

In order to determine whether there is any region of the coast where a marked change in the hydroid fauna occurs, the range of the tropical, temperate and endemic components was plotted separately (Fig. 3). (The range of species was used in preference to presence/absence, in order to minimize differences in collecting intensity.) This figure emphasizes the tropical nature of the east coast fauna, for there is a strong tropical component at the northern end which gradually drops away, until beyond St Helena Bay on the west coast (Sector 14) only one species remains for a short distance. The *rate* of disappearance is greatest at the boundary of Sectors 23/24, and is also high in the region of Sectors 13/14/15. The temperate and endemic components are both highest on the south coast, falling away fairly evenly to the north-east, but more sharply to the north-west in the region of Sectors 13/14/15. A similar trend appears when the range of species is plotted as a percentage of the total range (Fig. 4).

It is thus possible to distinguish two 'zones of change', though they are not as marked as might be expected. The first (Sectors 23/24) provides a convenient




Fig. 3. The distribution of the temperate component (above), the endemic component (centre) and the tropical component (below). From range of species.

boundary between the east and south coast fauna. It is located between 30 and 32°S latitude, i.e. the stretch of coast between the Bashee River in the south and Durban in the north. For the discussions which follow, the boundary between Sectors 23 and 24 is used (31°S). This is more or less in agreement with conclusions drawn from other faunal groups. Stephenson & Stephenson (1972) suggest a boundary zone between Port St Johns and Qolora, i.e. at about 32°S, for the intertidal fauna (the form of the graph of the tropical component in Figure 3 of this paper is very similar to that for the intertidal fauna in Stephenson & Stephenson (1972, fig. 8.26)); J. L. B. Smith (1949) and M.-L. Penrith (1970) suggest the Great Kei River at 32°42'S for the fishes, and Day (1967) suggests the Bashee River at 32°15'S for the benthic Polychaeta.




Fig. 4. The distribution of the temperate, endemic and tropical components, from range of species as a percentage of the total range. Sectors 1-7 have been omitted since the total number of species is less than twenty and the percentages unrealistic.

At the boundary of Sectors 23/24 the monthly minimum surface temperature near the coast does not fall below 16°C, the maximum does not rise above 27°C and the mean ranges from 21 to 24°C (U.S. Naval Oceanographic Office 1967). This is approximately the region where the Agulhas Bank begins to widen and force the warm Agulhas Current offshore. Further south the latter is separated from the coast by a cooler counter-current flowing northwards.

The usefulness of the second 'zone of change' in the south-westerly corner of the Cape Province (Sectors 13/14/15) is more debatable. It could be used to divide the temperate region, which stretches from north of 20°S on the west coast to 30–32°S on the east coast, into two provinces, if warranted by the fauna.

To test this an arbitrary boundary was placed at Cape Point (Sector 14/15) and the composition of species on the two sides of it compared with one another and with the east coast (Table 1). (Cape Point was used as a boundary by Stephenson (1947, fig. 3), Ekman (1953) and Briggs (1974).) It may be noted from Figures 3 and 4 that there is no evidence of a change in fauna at Cape Agulhas (Sector 15/16), the most southerly point of Africa).

TABLE 1

Analysis of fauna of west coast, as compared with that of the south and east coasts (actual records, not range).

Component	West coast (Sectors 1–14)		South coast (Sectors 15–23)		East coast (Sectors 24–36)	
	No. of species	%	No. of species	%	No. of species	%
Tropical	8	9,6	30	19,0	72	39,6
Temperate	13	15,7	20	12,7	12	6,6
Endemic to area	2	2,4	21	13,3	17	9,3
Endemic to more than one area .	25	30,1	41	25,9	30	16,5
Cosmopolitan	17	20,5	21	13,3	28	15,4
Scattered	18	21,7	25	15,8	23	12,6
Total	83	100,0	158	100,0	182	100,0

It is apparent from Table 1 that the west coast fauna consists mainly of endemic, cosmopolitan and scattered species. The tropical species diminish in number round the coast from east to west; only 8 species remain on the west coast and these are all absent north of St Helena Bay. The temperate species of the west coast are fewer in number than those of the south coast. Only 2 of them (*Clavopsella navis* and *Gonothyraea loveni*, both from Table Bay harbour and probably introduced by ships), and possibly a third (*Sarsia eximia*), are restricted to the west coast. Only 2 rare species are endemic solely to the west coast (*Myriothele tentaculata* and *Eudendrium ritchiei*), whereas 21 are endemic to the south coast and 17 to the east coast. Of the cosmopolitan and scattered components only 3 (*Acryptolaria crassicaulis*, *Hybocodon unicus* and *Hydrodendron gracilis*) have been recorded from the west coast alone.

In total, thus, the west coast has only 8 unique species (9,8%), of which only 2 (2,4%) are endemic to the area. Briggs (1974) requires at least 10 per cent endemic species for the designation of a separate province, and on this basis the west coast of South Africa does not qualify, at least for the hydroids. The hydroid fauna of the west coast seems to be mainly an impoverished south coast fauna.

Although Day (1967) accepted a separate west coast element for the littoral polychaets, he could find no marked difference between the south and west coasts for the sublittoral ones, and ascribed this to the fairly uniform water temperature on the bottom of the continental shelf (he gave figures of 12–14°C at 100 m from Port Elizabeth to Lüderitz, but the records collated by Hulley (1972) show a somewhat lower range of 8–13°C at 100 m between 20°S on the west coast and Port Elizabeth on the south coast). On the other hand there *is* a difference in the surface water temperature on the two sides of Cape Point (Day gives a range of 15–20°C for the south coast and 12–15°C for the west coast, but under certain conditions and for certain periods the south coast temperature may drop to as low as 11°C (during local upwelling) and rise as high as 24°C, while that of the west coast may drop to 9°C (M. J. Penrith 1976)). In this respect it might be noted that all hydroids are sublittoral in the sense that they cannot withstand drying and survive in the littoral zone only in damp overhangs or open runnels.

None of this really explains why there should be a diminution of hydroid species on the west coast and practically no endemic population of cold-temperate forms. Ekman (1953) associates the poverty of the benthos of the 'Namaqua coast' with the hydrogen sulphide resulting from frequent upwelling and red tides. Briggs (1974) suggests that certain species are prevented from spreading westwards round Cape Point by the absence of high summer temperatures for reproduction (he gives a February mean of 16–18°C for South West Africa as against one of 20–22°C for the south coast, but compare remarks in parenthesis above). M.-L. Penrith & Kensley (1970a) suggest that a region of minimal temperatures between Hondeklip Bay and Lüderitz (i.e. north of St Helena Bay) acts as a barrier to the establishment of southern warm-temperate intertidal species further north. The paucity of hydroid records on the west coast may be partly a reflection of the difficulty of collecting on the exposed and inhospitable Skeleton Coast. However, M.-L. Penrith (pers. comm.), who has collected in this area, feels that the poverty is a real one, and perhaps linked to the absence of suitable habitats such as sheltered coves, rock pools, etc., and the preponderance of surf beaches and sandy floors. The harsh conditions and poverty of the rocky-shore fauna are also emphasized by M. J. Penrith (1976).

There is no evidence at all of a southward spread of species from tropical west Africa. The southern boundary of the west African tropical region is too far north for consideration in this paper (14 or 15°S: Briggs 1974; 18–22°S: Hulley 1972). M.-L. Penrith & Kensley (1970b) found the first of the west African littoral species at Rocky Point (18°59'S).

In conclusion the south and west coast sublittoral areas are considered to represent a single temperate province for the hydroid fauna, which might be termed the Agulhas/Namaqua Province, the centre of the endemic population being on the Agulhas Bank.

THE TROPICAL COMPONENT

Of the total number of 251 species, 77 (30,7%) can be classified as tropical. Of these, 20 are circumtropical, 24 Indo-West-Pacific and 33 confined to the Western Indian Ocean. In addition there are 17 endemic species which are confined to the South African east coast, and therefore with tropical tendencies (these were termed 'subtropical endemics' by Day (1974)). (Appendix 2.)

Ekman (1953) and Briggs (1974) consider that the main centre of origin of the tropical fauna is the Indo-Malayan Region. From here species tend to spread west to the Indian Ocean (and may penetrate into the Red Sea or round southern Africa), and to a lesser extent east to the Eastern Pacific.

Species have been classified here as Indo-West-Pacific or Western Indian Ocean even though a few have one or two records outside these areas. Thus, 2 have penetrated into the Mediterranean (*Corydendrium parasiticum*, *Thyroscyphus fruticosus*), 4 have reached the tropical west coast of Africa (*Abietinaria laevimarginata*, *Campanularia africana*, *Corydendrium parasiticum*, *Sertularia ligulata*), 1 has reached tropical South America (*Dentitheca bidentata*), and 1 has reached southern Australia (*Dynamena obliqua*). Further spread would produce a circumtropical distribution, but of the species classified as such no less than 13 have failed to reach the east coast of the Pacific, and this applies also to 4 of the cosmopolitans (Table 2). There is therefore support for Briggs's contention (1974) that the 'East Pacific barrier' is one not easily crossed.

There is some evidence for a secondary centre of origin in the Western Indian Ocean, for in addition to the seventeen South African subtropical endemics (i.e. present on the east coast only), there are seven Western Indian

TABLE 2

List of circumtropical and cosmopolitan species which do not occur on the east coast of the Pacific.

Circumtropical	Cosmopolitan
<i>Diphasia digitalis</i>	<i>Antennella secundaria</i>
<i>Dynamena quadridentata</i>	<i>Cladocoryne floccosa</i>
<i>Gymnangium hians</i>	<i>Coryne pusilla</i>
<i>Halecium dyssymetrum</i>	<i>Halecium sessile</i>
<i>Hydrodendron caciniformis</i>	
<i>Idiellana pristis</i>	
<i>Monostaechas quadridens</i>	
<i>Plumularia strictocarpa</i>	
<i>Scandia mutabilis</i>	
<i>Sertularella diaphana</i>	
<i>Sertularia distans</i>	
<i>S. marginata</i>	
<i>S. turbinata</i>	

Ocean species which, though absent from the Indo-Polynesian area, have yet penetrated into the Red Sea (*Clytia latithea*, *Cytaeis nassa*, *Diphasia heurteli*, *Halopteris glutinosa*, *Hydractini kaffraria*, *Plumularia wasini*, *Solanderia minima*).

Some communication between southern Australia and the Western Indian Ocean is indicated, since there are five species common to the two, namely *Amphisbetia maplestoni*, *Craterithea acanthostoma*, *Halopteris glutinosa*, *Sertularella arbuscula* and *Synthecium dentigerum*. These occur nowhere else (except for *H. glutinosa* which also occurs in the Red Sea) and the likelihood is that the movement has been from west to east with the West Wind Drift.

THE TEMPERATE COMPONENT

The total temperate component includes 28 species (11,2%), of which 11 are antitropical (bipolar), 10 are restricted to the Southern hemisphere and 7 are restricted to the Atlantic (Appendix 2). These are species which cannot survive the high temperatures of the tropics, although some can spread into the cooler waters of the subpolar regions. The southern species must have evolved in the south, but the antitropical and Atlantic species may have evolved in either the south or the north and must at some time have crossed the tropics. This may have been achieved by equatorial submergence, for Briggs (1974) points out that the tropical surface water is only about 30–40 m deep in the Eastern Atlantic. It may also be due to transportation on the hulls of ships. In all, twenty-one species have been recorded on ships' hulls or floating objects such as buoys. The fauna of ships' hulls is a characteristic group of hardy species (Millard 1952), most of them cosmopolitan. Three of the species (all belonging to the temperate component) are known only in harbour areas in South Africa, and it is suggested that these are recent introductions by ships:

Gonothyraea loveni is well established in Cape Town docks, occurring abundantly on pylons, cables, etc., as well as on hulls of ships. It has not been reported from any other part of South Africa. Elsewhere it is known from the North Atlantic, New Zealand, Tasmania and the Argentine.

Campanularia calceolifera has been found, and reproducing, in Cape Town docks on a barge and on a raft, neither of which had left the harbour. It must, therefore, be established, though not common. It is not known for certain elsewhere in South Africa, a record from False Bay by Stechow (1925) being of infertile material and thus subject to doubt. The species is well known on European and Mediterranean coasts and also occurs on the east coast of North America.

Clavopsella navis is known from a ship's hull in Cape Town docks, and also from the Kiel Canal (Thiel 1962 as *C. quadranularia*). The latter is probably the original locality since the colonies there were more abundant and better developed.

Since all these three species occur both in the South and North Atlantic they must have traversed the tropics, though they have not established themselves in tropical latitudes so far as is known. They may have been hardy enough to survive a brief passage through warmer waters on ships' hulls, or they may have died back during transport, leaving the hydrorhiza to regenerate again in suitable temperature conditions. Ralph (1961) also found evidence of the introduction of hydroids by ships' hulls to New Zealand harbours.

It is interesting to compare the relations of the temperate species with the other two southern continents. From Table 3 it can be seen that South Africa has 9 species in common with southern Australasia alone, but only 3 species in common with South America alone. There are 7 species common to all three.

TABLE 3

Temperate species common to the southern continents.

South Africa and South Australasia	South Africa and temperate South America	Temperate part of all three southern continents
<i>Kirchenpaueria triangulata</i>	<i>Phialella turrita</i>	<i>Amphisbetia minima</i>
<i>Nemertesia ciliata</i>	<i>Sertularella striata</i>	<i>Filellum antarcticum</i>
<i>Parascyphus simplex</i>	<i>Symplectoscyphus paulensis</i>	<i>Gonothyrea loveni</i>
<i>Plumularia obliqua</i>		<i>Nemertesia cymodocea</i>
<i>P. spinulosa</i>		<i>Plumularia filicaulis</i>
<i>Sarsia eximia</i>		<i>Staurocladia vallentini</i>
<i>Sertularella annulaventricosa</i>		<i>Tubularia larynx</i>
<i>Stereotheca elongata</i>		
<i>Symplectoscyphus macrogonus</i>		
(Total: 9)	(Total: 3)	(Total: 7)

This would suggest dispersal eastward from the Agulhas Bank. Beurois (1975) emphasized the role of eastward drifting kelp in the dispersal of attached forms to St Paul and Amsterdam Islands, and Arnaud *et al.* (1976) discussed the transport of South African fauna to St Helena on kelp. If the list in Table 3 is extended to include the cosmopolitan and scattered components and those tropical forms which have spread south of the Tropic of Capricorn, the relationship between South Africa and Australasia is more marked (34 species in common with Australasia alone, 10 with South America alone, and 26 in all three).

The distribution of those South African species which have antarctic or subantarctic affinities is given in Table 4. In all, 41 species are listed, a relatively small number (16,3% of the total), and of these half (21) are cosmopolitan and this tends to mask the relationships. Without the cosmopolitan species only 20 (8,0%) have antarctic affinities and of these 11 are classified as temperate. The absence from South Africa of certain common antarctic genera, such as *Staurotheca*, *Tulpa*, *Grammaria* and *Silicularia*, is very striking. On the whole there is very little evidence of relationship with the Antarctic.

TABLE 4

Distribution of species with antarctic/subantarctic affinities. The Magellan Region includes Gauss Station, the Falklands and South Georgia. The Kerguelen Group includes Marion, Prince Edward and Crozet Islands.

Species	Component	Antarctic continent and neighbourhood	Magellan Region	Tristan group	Kerguelen group	St Paul
<i>Acryptolaria conferta</i>	Cosmopolitan	×	×			
<i>A. crassicaulis</i>	Scattered	×	×			
<i>Aglaophenia p. pluma</i>	Cosmopolitan		×			×
<i>Amphisbetia minima</i>	Temperate (Antitr.)		×			×
<i>A. operculata</i>	Cosmopolitan		×		×	×
<i>Antennella quadriaurita</i>	Scattered			×		
<i>A. secundaria</i>	Cosmopolitan					×
<i>Bimeria vestita</i>	Cosmopolitan		×			
<i>Bougainvillia macloviana</i>	Scattered		×		×	
<i>Campanularia crenata</i>	Cosmopolitan		×			
<i>C. integra</i>	Cosmopolitan		×			
<i>Clytia hemisphaerica</i>	Cosmopolitan		×			×
<i>C. paulensis</i>	Scattered	×				×
<i>Coryne pusilla</i>	Cosmopolitan				×	
<i>Dynamena cornicina</i>	Cosmopolitan					×
<i>Filellum antarcticum</i>	Temperate (Southern)	×	×			
<i>F. serratum</i>	Cosmopolitan		×		×	
<i>Halecium beanii</i>	Cosmopolitan		×			
<i>H. delicatulum</i>	Cosmopolitan	×	×		×	
<i>H. tenellum</i>	Cosmopolitan	×	×	×	×	
<i>Hincksella echinocarpa</i>	Temperate (Southern)				×	
<i>Hybocodon unicus</i>	Scattered		×			
<i>Kirchenpaueria triangulata</i>	Temperate (Southern)				×	
<i>Lafoea benthophila</i>	Scattered	×	×		×	×
<i>L. dumosa</i>	Cosmopolitan	×	×		×	
<i>Modeeria rotunda</i>	Cosmopolitan		×		×	
<i>Nemertesia cymodocea</i>	Temperate (Southern)		×			
<i>N. ramosa</i>	Scattered		×			
<i>Obelia bidentata</i>	Cosmopolitan		×	×	×	
<i>O. dichotoma</i>	Cosmopolitan	×	×	×		×
<i>O. geniculata</i>	Cosmopolitan		×	×	×	×
<i>Parascyphus simplex</i>	Temperate (Antitr.)	×		×		
<i>Phialella turrita</i>	Temperate (Antitr.)		×			
<i>Plumularia pulchella</i>	Scattered		×			
<i>P. setacea</i>	Cosmopolitan		×			×
<i>Sertularella leiocarpa</i>	Temperate (Southern)			×		×
<i>S. p. polyzonias</i>	Cosmopolitan		×		×	
<i>S. striata</i>	Temperate (Southern)		×			
<i>Staurocladia vallentini</i>	Temperate (Southern)		×		×	
<i>Symplectoscyphus paulensis</i>	Temperate (Southern)		×			×
<i>Zygophylax armata</i>	Scattered			×		

THE ENDEMIC COMPONENT

The number of endemic species strictly belonging to the South African coast south of 20°S is 76. However, there are 2 species which also occur immediately north of 20° on the west coast (*Gattya humilis* and *Salacia articulata*). 4 species occur only in South Africa and the Vema Seamount (*Halopteris pseudoconstricta*, *Salacia articulata*, *Sertularella flabellum* and *S. megista*), and it was shown by Millard (1966) that the hydroid fauna of the Vema Seamount is essentially South African (21 of the 23 species also occur in South Africa). If these 6 species are included, the endemics number 82 (32,7% of the total of 251). It might be noted, however, that Mergner & Wedler (1977) have recorded *Eudendrium deciduum* and *Sertularella natalensis* from the Red Sea with a query.

As a comparison the following percentages of endemics have been given for other large marine groups in southern Africa. The figures are not strictly comparable, however, since the boundaries used for the 'Southern African Region' are not always the same.

Polychaeta	36 %	(Day 1967)
Echinoderms, other than Holothuria	48 %	(Clark & Courtman-Stock 1976)
Ascidians	c. 50 %	(Millar 1971)
Fish	25,3 %	(M. M. Smith 1970: 380 out of 1 500 species)

The south coast of southern Africa has more endemic species of hydroid than any other region, and the Agulhas Bank seems to be the centre of distribution of the endemic population (Figs 3–5).

There are only three genera endemic to South Africa, *Bicorona*, *Hydrocorella* and *Uniscyphus*, and all of these are monotypic. However, certain genera have a larger proportion of endemic species than others (Table 5). These are probably genera in which speciation is actively occurring in the South African area. They are also genera in which many of the species are not easily distinguished from one another, and thus pose constant problems to the systematist. Similarly, species with several subspecies are ones likely to be undergoing active speciation. Thus, *Thecocarpus flexuosus* has four subspecies; the nominal one is

TABLE 5

List of South African hydroid genera with the largest number of endemic species and/or subspecies.

Genus	Number of endemic species or subspecies	Total number of species or subspecies in South Africa
<i>Cladocarpus</i>	9	13
<i>Corhiza</i>	5	5
<i>Halopteris</i>	4	6
<i>Hydractinia</i>	3	5
<i>Sertularella</i>	14	23
<i>Zygophylax</i>	4	7




Fig. 5. An analysis of the distribution of the endemic species, with the total number of species distributed between six components, as in Stephenson (1972). From range of species.

tropical Indo-West-Pacific and the other three are endemic to southern Africa. *Sertularella polyzonias* also has four subspecies; the nominal one is cosmopolitan and two of the others are endemic to southern Africa.

Several common genera, on the other hand, have no southern African endemics: *Aglaophenia*, *Amphisbetia*, *Dynamena*, *Lytocarpus*, *Nemertesia*, *Obelia* and *Sertularia*.

THE COSMOPOLITAN AND SCATTERED COMPONENTS

The cosmopolitan species number 28 (11,2%). They owe their wide distribution to the fact that they are eurythermic, hardy and adaptable. The wide distribution possibly indicates a very old history, in which there has been plenty of time to disperse.

Species classified as scattered or disjunct number 36 (14,3%). Most of them

have odd distributions which do not fit any of the other categories. There are two possibilities. Either these are newly evolved eurythermic species which are in the process of spreading rapidly and likely to become cosmopolitan, or they are 'peripheral relicts' of fairly primitive forms which were once more widely distributed.

Among the scattered species three are interesting because outside southern Africa they are limited to the west and east coasts of the tropical Atlantic (*Clytia hummelincki*, *Zyzzyzus solitarius*) or to the western tropical Atlantic only (*Cladocarpus tenuis*). Three others have a similar distribution but also extend further north and/or east (*Aglaophenia latecarinata*, *A. pluma dichotoma*, *Symplectoscyphus amphoriferus*).

Cosmopolitan and scattered species with antarctic/subantarctic affinities are included in Table 4.

DISTRIBUTION AND DEPTH

Figure 6 shows that there are more species present in the shallow water (0–100 m) than at any other level, and although there have been fewer samples from deep water it is not expected that more extensive collecting will alter the picture to any great extent. The shallow water is obviously the optimum zone for hydroids. Most of the species in the littoral area are at the upper edge of




Fig. 6. The distribution of species according to depth. From range of species.

their depth range and able to survive only under overhangs and in rock pools and crevices. Only 24 of the 98 littoral species have not been found below low water of springs.

Deep-water species are those which occur over the edge of the continental shelf, and Ekman (1953) places an arbitrary upper boundary for the deep-sea fauna at 400 m. The number of hydroids known from below this depth round the South African coast is relatively small ($45 = 17.9\%$), and only eight of these reach 1 000 m (abyssal). The species are listed in Table 6.

These species can be divided into two arbitrary categories on the basis of their depth range both in South Africa and the rest of the world:

Stenobathic: those that are restricted to water deeper than 200 m.

Eurybathic: those that occur above and below 200 m.

These two categories are distributed between the different geographical components as follows:

	<i>Stenobathic</i>	<i>Eurybathic</i>	<i>Total</i>
Tropical	5	5	10
Temperate	1	4	5
Endemic	5	6	11
Cosmopolitan	0	9	9
Scattered	1	9	10
Total	12	33	45

All the (12) stenobathic species are in reality cold-water species frequenting water below about 16°C (Ekman 1953, table 42), and this includes five so-called 'tropical' species. These species are thus also stenothermic. Their limiting factor for distribution may be depth or temperature or a combination of both.

Some of the eurybathic species are also stenothermic. This applies to the following tropical and scattered species whose records *within the tropics* are never shallow, and which are thus restricted to cold water:

	<i>Most shallow record in tropics (m)</i>
<i>Acryptolaria rectangularis</i> (tropical)	228
<i>Cladocarpus tenuis</i> (scattered)	185
<i>Cryptolarella abyssicola</i> (scattered)	4 560
<i>Symplectoscyphus amphoriferus</i> (scattered)	161
<i>Zygophylax sibogae</i> (scattered)	Only one tropical record: depth not given.

To the same category of eurybathic/stenothermic species might be added the endemic and temperate eurybathic species, at any rate those restricted to the south-west coast of South Africa and the east coast below 200 m:

Cladocarpus sinuosus (endemic): minimum east coast record 400 m.

Hincksella echinocarpa (temperate): minimum record Kerguelen 41 m, minimum east coast record 1 610 m.

TABLE 6

South African deep-water species.

SB—stenobathic (never recorded above 200 m); EB—eurybathic (occurring at all depths);
ST—stenothermic (cold-water species); ET—eurythermic (occurring at all temperatures).

Species	Component	Depth range in South Africa (m)	Depth range outside South Africa (m)	Component of depth/ temperature
<i>Acryptolaria conferta</i>	Cosmopolitan	64–1200	0–4400	EB/ET
<i>A. crassicaulis</i>	Scattered	2835	109–767	EB/ET
<i>A. rectangularis</i>	Tropical	110–1200	228–600	EB/ST
<i>Antennella quadriaurita</i>	Scattered	0–825	0–400	EB/ET
<i>Branchiocerianthus imperator</i>	Tropical	730	200–5290	SB/ST
<i>Campanularia hincksii</i>	Cosmopolitan	9–450	0–811	EB/ET
<i>Cladocarpus distomus</i>	Tropical	292–2200	55–1362	EB/ET
<i>C. dofleini</i>	Tropical	425–550	1019	SB/ST
<i>C. inflatus</i>	Endemic	495	—	SB/ST
<i>C. millardae</i>	Tropical	200–595	5020	SB/ST
<i>C. natalensis</i>	Endemic	400–900	—	SB/ST
<i>C. sinuosus</i>	Endemic	183–680	—	EB/ST
<i>C. tenuis</i>	Scattered	495	185–550	EB/ST
<i>Clytia gravieri</i>	Tropical	0–528	0–4243	EB/ET
<i>Cryptolarella abyssicola</i>	Scattered	100–2740	745–4970	EB/ST
<i>Filellum serratum</i>	Cosmopolitan	24–900	0–650	EB/ET
<i>Garveia crassa</i>	Tropical	625–900	741	SB/ST
<i>Halecium beanii</i>	Cosmopolitan	0–780	0–1134	EB/ET
<i>H. sessile</i>	Cosmopolitan	425–430	0–752	EB/ET
<i>H. tenellum</i>	Cosmopolitan	0–900	0–730	EB/ET
<i>Halopteris glutinosa</i>	Tropical	0–550	1–80	EB/ET
<i>H. polymorpha</i>	Tropical	0–900	0–89	EB/ET
<i>H. tuba</i>	Endemic	11–550	—	EB/ET
<i>Hincksella echinocarpa</i>	Temperate	1610–2200	41–109	EB/ST
<i>Hydrocorella africana</i>	Endemic	0–500	—	EB/ET
<i>Kirchenpaueria triangulata</i>	Temperate	111–1207	550–585	EB/ST
<i>Lafoea benthophila</i>	Scattered	425–430	672–3237	SB/ST
<i>L. dumosa</i>	Cosmopolitan	60–920	0–2200	EB/ET
<i>Modeeria rotunda</i>	Cosmopolitan	70–550	0–1240	EB/ET
<i>Nemertesia antennina</i>	Scattered	425–430	0–1779	EB/ET
<i>N. ramosa</i>	Scattered	11–700	0–872	EB/ET
<i>Plumularia antonbruuni</i>	Endemic	440	—	SB/ST
<i>P. mossambicae</i>	Endemic	110–550	—	EB/ET
<i>P. setacea</i>	Cosmopolitan	0–430	0–752	EB/ET
<i>Sertularella leiocarpa</i>	Temperate	200–900	183–672	EB/ST
<i>Stegolaria geniculata</i>	Tropical	775–850	253–910	SB/ST
<i>Symplectoscyphus amphoriferus</i>	Scattered	550–900	161–1256	EB/ST
<i>S. arboriformis</i>	Endemic	10–420	—	EB/ET
<i>S. paulensis</i>	Temperate	347–1200	400–672	SB/ST
<i>Uniscyphus fragilis</i>	Endemic	360–420	—	SB/ST
<i>Zygophylax africana</i>	Endemic	137–850	—	EB/ST
<i>Z. armata</i>	Scattered	48–440	183	EB/ET
<i>Z. brownei</i>	Temperate	400–550	20–752	EB/ET
<i>Z. inconstans</i>	Endemic	360–450	—	SB/ST
<i>Z. sibogae</i>	Scattered	88–900	146–550	EB/ST

Summary: Cosmopolitan . . . 9
 Scattered . . . 10
 Tropical . . . 10
 Temperate . . . 5
 Endemic . . . 11

Total . . . 45

Kirchenpaueria triangulata (temperate): minimum record Agulhas Bank 111 m, minimum east coast record 350 m.

Sertularella leiocarpa (temperate): minimum record Tristan 183 m, minimum east coast record 280 m.

Zygophylax africana (endemic): minimum east coast record 400 m.

For the (10) species in this eurybathic/stenothermic category the limiting factor is obviously temperature rather than depth.

The remaining (23) species are truly eurybathic and eurythermic, for they can occur in any temperature and at any depth. As might be expected, the majority (13) are cosmopolitan or scattered.

Hedgpeth (1957) suggested that all the deep-sea benthic population will eventually prove to be cosmopolitan, but it is clear from this analysis that geographically cosmopolitan species are not necessarily eurythermic.

The data for a selected group of 21 species were examined to see whether the depth distribution round the coast bore any relation to changing water temperatures. These were the species recorded from a minimum of 10 sectors and from depths extending below 100 m. Only 7 of the species so tested showed indications of an increasing minimum depth as one passes round the coast from west to east and then north; only 5 showed indications of an increasing maximum depth (Fig. 7, Table 7). This might be interpreted as an inability in a few species to survive in the warmer surface and shallow waters of the east coast. It is not surprising that none of these species belongs to the tropical component.

Cladocarpus is a characteristic deep-water genus, and 12 of the 13 South African species extend down below 100 m, 9 below 200 m, 3 below 500 m and 7 do not occur in water shallower than 200 m.

TABLE 7

The relationship between depth and distribution.

A. The minimum recorded depth in metres; B. The maximum recorded depth in metres.

Species	Sectors							Component
	0-5	6-10	11-15	16-20	21-24	25-30	31-35	
A. <i>Antennella quadriaurita</i>	—	35	0	9	50	100	—	Scattered
<i>Halecium delicatulum</i>	—	—	0	4	27	110	55	Cosmopolitan
<i>Halopteris tuba</i>	—	—	22	11	27	70	110	Endemic
<i>Salacia articulata</i>	0	23	0	10	0	18	110	Endemic
<i>Sertularella flabellum</i>	—	—	15	10	27	124	—	Endemic
<i>Sertularella megista</i>	—	—	0	10	27	49	110	Endemic
<i>Symplectoscyphus arboriformis</i>	0	10	11	11	49	124	—	Endemic
B. <i>Antennella quadriaurita</i>	—	40	79	114	210	825	—	Scattered
<i>Salacia articulata</i>	0	23	75	110	91	135	132	Endemic
<i>Sertularella flabellum</i>	—	—	99	200	91	232	—	Endemic
<i>Sertularella megista</i>	—	—	75	200	164	219	347	Endemic
<i>Symplectoscyphus arboriformis</i>	—	20	35	120	90	420	—	Endemic




Fig. 7. The distribution according to depth of three species which show tendencies towards greater depths in warmer waters. Actual records.

ESTUARINE AND BRACK-WATER SPECIES

Although many hydroids penetrate into the mouths of estuaries, especially on rocky substrata, only two species are truly estuarine and have not been found in the open sea. These are *Bimeria fluminalis*, which grows on mangrove roots in Richards Bay and St Lucia estuaries, and *Hydractinia kaffraria*, which is a commensal on the shell of *Nassa kraussiana* and occurs in estuaries from the Breede River mouth on the south coast to Durban on the east coast.

The Limnomedusan, *Ostroumovia inkermanica*, is known from several brack-water lakes on the east coast, the hydroid from Lagoa Poelala, Mozambique, in a salinity of 8 parts per thousand, both hydroid and medusa from Nhlanga Lake, Zululand, in 3–4 parts per thousand, and the medusa alone from




Fig. 8A-D. Distribution of six representative endemic species: A. *Lovenella chiquitita* (west coast + False Bay), *Cladocarpus leloupi* (south coast) and *Sertularella natalensis* (east coast). B. *Corhiza scotiae* (west + south coast). C. *Syntheicum hians* (south + east coast). D. *Salacia articulata* (west + south + east coast). E-F. Distribution of two representative temperate species: E. *Symplectoscyphus macrogonus* (west + south coast). F. *Sertularella leiocarpa* (west + south + east coast), also stenothermic.




Fig. 9A-D. Distribution of four representative tropical species from east coast: A. *Lytocarpus philippinus* (to just north of Durban). B. *Dynamena crisioides* (to East London). C. *Sertularia turbinata* (to Mossel Bay). D. *Sertularella arbuscula* (to Saldanha Bay). E-F. Distribution of two representative cosmopolitan species: E. *Hebella scandens* (St Helena Bay to Mozambique). F. *Amphisbetia operculata* (Lüderitz to Natal).

Lake St Lucia North in 10,4–10,8 parts per thousand.

CONCLUSION

The composition of the southern African hydroid fauna is summarized in Table 8, whence it can be seen that there is a mixture of endemic, tropical, temperate and cosmopolitan species in that order of abundance.

The presence of a large endemic population, the particular distribution of some of the Western Indian Ocean tropical species (p. 169), and the apparent proliferation of subspecies in certain genera (p. 173), all point to an active evolutionary centre in the South African area.

It is clear that there has also been a heavy invasion of tropical species from the Indo-West-Pacific region, and these species form the major component of the fauna of the east coast.

Regarding the temperate and cosmopolitan species, and considering

- (i) the low relationship of the South African temperate fauna to that of the antarctic and subantarctic regions on the one hand, and to that of South America and Australasia on the other,
- (ii) the practically certain role of ships in the introduction of at least three species out of 251 (1,2%) to South Africa within a period of some 300 years (p. 170),

TABLE 8

Summary of analysis of geographical components of South African hydroid fauna (from actual records, not range).

<i>Component</i>	<i>West-plus-south coast: Sectors 1–23 (171 species)</i>		<i>East coast: Sectors 24–34 (182 species)</i>		<i>Whole of South Africa: Sectors 1–34 (251 species)</i>	
	No.	Percentage	No.	Percentage	No.	Percentage
TROPICAL						
Circumtropical . . .	10	5,8	18	9,9	20	8,0
Indo-West-Pacific . . .	7	4,1	23	12,6	24	9,6
W. Indian Ocean . . .	16	9,4	31	17,0	33	13,1
		19,3		39,6		30,7
TEMPERATE						
Antitropical	10	5,8	5	2,7	11	4,4
Southern	8	4,7	5	2,7	10	4,0
Atlantic	5	2,9	2	1,1	7	2,8
		13,4		6,5		11,2
ENDEMIC						
To one area only . . .	35	20,5	17	9,3	52	20,7
To the whole of South Africa	30	17,5	30	16,5	30	12,0
		38,0		25,8		32,7
UNCLASSIFIED						
Cosmopolitan	22	12,9	28	15,4	28	11,2
Scattered	28	16,4	23	12,6	36	14,3
		29,3		28,0		25,5

(iii) the fact that drifting weed and flotsam has probably performed a similar role throughout the ages, it is hardly necessary to evoke the Theory of Continental Drift to explain the distribution.

REFERENCES

- ARNAUD, F., ARNAUD, P. M., INTÈS, A. & LE LOEUFF, P. 1976. Transport d'Invertébrés benthiques entre l'Afrique du Sud et Sainte Hélène par les lamineuses (Phaeophyceae). *Bull. Mus. natn. Hist. nat. Paris* (3) **384**, Écol. gén. **30**: 49–55.
- BEUROIS, J. 1975. Étude écologique et halieutique des fonds de pêche et des espèces d'intérêt commercial (langoustes et poissons) des îles Saint-Paul et Amsterdam (Océan Indien). *CNFR* **37**: 1–91.
- BLANCO, O. M. 1967. Contribucion al conocimiento de los hidrozoarios Argentinos. *Revta Mus. La Plata* (N.S.) Zool. **9**: 243–297.
- BLANCO, O. M. 1968. Nueva contribucion al conocimiento de la fauna marina Hidroide. *Revta Mus. La Plata* (N.S.) Zool. **10**: 195–224.
- BLANCO, O. M. 1973. Nuevos plumularidos para aguas Argentinas. *Neotropica* **19**: 73–78.
- BLANCO, O. M. 1974. Adicion a los hidrozoos Argentinos. *Neotropica* **20**: 40–47.
- BLANCO, O. M. & BELLUSCI DE MIRALLES, D. A. 1972a. Nuevos aportes a los campanularidos de Argentina. *Revta Mus. La Plata* (N.S.) Zool. **11**: 137–151.
- BLANCO, O. M. & BELLUSCI DE MIRALLES, D. A. 1972b. Hidrozoos de la Isla Pedro I. *Contrnes Inst. antart. argent.* **145**: 3–43.
- BRIGGS, J. C. 1974. *Marine zoogeography*. New York, etc.: McGraw-Hill.
- BUCHANAN, J. B. 1957. The hydroid fauna of the Gold Coast. *Revue Zool. Bot. afr.* **56**: 349–372.
- CALDER, D. R. 1975. Biotic census of Cape Cod Bay: hydroids. *Biol. Bull. mar. biol. Lab. Woods Hole* **149**: 287–315.
- CALDER, D. R. 1976. The zonation of hydroids along salinity gradients in South Carolina. In: *Coelenterate ecology and behaviour*, ed. G. O. Mackie: 165–174. New York & London: Plenum Press.
- CLARK, A. M. & COURTMAN-STOCK, J. 1976. *The echinoderms of southern Africa*. London: British Museum.
- COOKE, W. J. 1975. Shallow water hydroids from Enewetak Atoll, Marshall Islands. *Micronesica* **11**: 85–108.
- CORNELIUS, P. F. S. 1975a. The hydroid species of *Obelia* (Coelenterata, Hydrozoa: Campanulariidae), with notes on the medusa stage. *Bull. Br. Mus. nat. Hist. (Zool.)* **28**: 251–292.
- CORNELIUS, P. F. S. 1975b. A revision of the species of Lafoeidae and Haleciidae (Coelenterata: Hydroida) recorded from Britain and nearby seas. *Bull. Br. Mus. nat. Hist. (Zool.)* **28**: 375–426.
- DAY, J. H. 1967. *A monograph on the Polychaeta of southern Africa*. London: British Museum.
- DAY, J. H. 1974. The ecology of Morrumbene Estuary, Moçambique. *Trans. R. Soc. S. Afr.* **41**: 43–97.
- EKMANN, S. 1953. *Zoogeography of the sea*. London: Sidgwick & Jackson.
- HEDGPETH, J. W. 1957. Marine biogeography. In: *Treatise on marine ecology and paleoecology*. I. *Mem. geol. Soc. Am.* **67**: 359–382.
- HULLEY, P. A. 1972. The origin, interrelationships and distribution of southern African Rajidae (Chondrichthyes, Batoidei). *Ann. S. Afr. Mus.* **60**: 1–103.
- MERGNER, H. & WEDLER, E. 1977. Über die Hydroidpolypenfauna des Roten Meeres und seiner Ausgänge. *Meteor ForschErgebn. (D)* **24**: 1–32.
- MICHEL, C. 1974. Notes on marine biology studies made in Mauritius. *Bull. Maur. Inst.* **7** (2): 1–284.
- MILLAR, R. H. 1971. The biology of ascidians. *Adv. mar. Biol.* **9**: 1–100.
- MILLARD, N. A. H. 1952. Observations and experiments on fouling organisms in Table Bay harbour, South Africa. *Trans. R. Soc. S. Afr.* **33**: 415–445.
- MILLARD, N. A. H. 1966. Hydroids of the Vema Seamount. *Ann. S. Afr. Mus.* **48**: 489–496.
- MILLARD, N. A. H. 1975. Monograph on the Hydroida of southern Africa. *Ann. S. Afr. Mus.* **68**: 1–513.

- MILLARD, N. A. H. 1977a. Hydroids from the Kerguelen and Crozet shelves, collected by the cruise MD.03 of the *Marion-Dufresne*. *Ann. S. Afr. Mus.* **73**: 1-47.
- MILLARD, N. A. H. 1977b. The South African Museum's *Meiring Naude* cruises. Part 3. Hydroida. *Ann. S. Afr. Mus.* **73**: 105-131.
- MILLARD, N. A. H. & BOUILLON, J. 1975. Additional hydroids from the Seychelles. *Ann. S. Afr. Mus.* **69**: 1-15.
- PENRITH, M. J. 1976. Distribution of shallow water marine fishes around southern Africa. *Cimbebasia* (A) **4**: 137-154.
- PENRITH, M.-L. 1970. The distribution of the fishes of the family Clinidae in southern Africa. *Ann. S. Afr. Mus.* **55**: 135-150.
- PENRITH, M.-L. & KENSLEY, B. F. 1970a. The constitution of the intertidal fauna of rocky shores of South West Africa. Part I. — Lüderitzbucht. *Cimbebasia* (A) **1**: 191-239.
- PENRITH, M.-L. & KENSLEY, B. 1970b. The constitution of the fauna of rocky intertidal shores of South West Africa. Part II. — Rocky Point. *Cimbebasia* (A) **1**: 243-268.
- RALPH, P. M. 1961. New Zealand thecate hydroids. Part V. — The distribution of the New Zealand thecate hydroids. *Trans. R. Soc. N. Z. (Zool.)* **1**: 103-111.
- SMITH, J. L. B. 1949. *The sea fishes of southern Africa*. Cape Town: Central News Agency.
- SMITH, M. M. 1970. Endemism in South African fishes. In: *Oceanography in South Africa*. Mimeographed Res. SANCOR Sympos., Durban, No. H3: 1-11.
- STECHOW, E. 1925. Hydroiden der deutschen Tiefsee-Expedition. *Wiss. Ergebn. dt. Tiefsee-Exped. 'Valdivia'* **17**: 383-546.
- STEPHENSON, T. A. 1947. The constitution of the intertidal fauna and flora of South Africa. Part III. *Ann. Natal Mus.* **11**: 207-324.
- STEPHENSON, T. A. & STEPHENSON, A. 1972. *Life between tidemarks on rocky shores*. San Francisco: W. H. Freeman & Co.
- THIEL, H. 1962. *Clavopsella quadranularia* nov. spec. (Clavopsellidae nov. fam.), ein neuer Hydroidpolyp aus der Ostsee und seine phylogenetische Bedeutung. *Z. morph. Ökol. Tiere* **51**: 227-260.
- U.S. NAVAL OCEANOGRAPHIC OFFICE. 1967. Monthly charts of mean, minimum, and maximum sea surface temperature of the Indian Ocean. *Spec. Publ. nav. oceanogr. Off., Washington* **99**: 1-48.
- WATSON, J. E. 1975. Hydroids of Bruny Island, southern Tasmania. *Trans. R. Soc. S. Aust.* **99**: 157-176.

APPENDIX 1

List of new hydroid records identified by the author since the publication of Millard (1975, 1977*a*, 1977*b*) and incorporated in this paper.

FOREIGN (South African species recorded outside the country)		Locality data	SAM no.
Family and species			
Solanderiidae			
<i>Solanderia minima</i> (Hickson, 1903)		Tulear, Madagascar; submitted by J. Bouillon, 1976	
Lafoeidae			
<i>Filellum serratum</i> (Clarke, 1879)		Vema Seamount; coll. J. R. Grindley	H 1653
<i>Lafaea dumosa</i> (Fleming, 1820)		South Georgia; coll. P. A. Hulley	H 2912
Campanulariidae			
<i>Obelia bidentata</i> Clarke, 1875		Nightingale Island; coll. Tristan Investments	H 1990
Plumulariidae			
<i>Aglaophenia pluma parvula</i> Bale, 1882		Gough Island, dredged; coll. Tristan Investments	H 1948
<i>Antennella quadriaurita</i> Ritchie, 1909		Nightingale Island; Tristan; coll. Tristan Investments	H 1991, 1949
SOUTH AFRICAN (N.B. Records are taken as 'new' if they have not been recorded before from the particular latitude/longitude square)			
* — new record for South Africa. MD — from <i>Meiring Naude</i> , Cruise 3. Depths: l — littoral; s — 1 to 99 m; md — 100 to 199 m; d — 200 to 499 m; vd — 500 to 999 m; a — 1 000 + m.			
Family and species		Locality data, including S — latitude/E — longitude square	SAM no.
Bougainvilliidae			
<i>Garveia crassa</i> (Stechow, 1923)		Off Natal: 30/30 (vd); MD	H 2933
Pandeidae			
<i>Hydrichthys boycei</i> Warren, 1916		Eastern Cape: 33/27 (s), 33/26; on <i>Acanthurus lineolatus</i> and <i>Chirodactylus brachydactylus</i> ; coll. R. Winterbottom	H 1970, 1985
Campanulinidae			
<i>Stegolaria geniculata</i> (Allman, 1888)		Off Natal: 30/30 (vd); MD	H 2943
Haleciidae			
<i>Halecium beanii</i> (Johnston, 1838)		Sodwana Reef: 27/32 (s); coll. A. E. Louw. Off Natal: 30/30 (vd); MD	H 2935, 2897
<i>H. delicatulum</i> Coughtrey, 1876		Jeffreys Bay: 34/24; coll. Mrs C. Giles	
<i>H. dichotomum</i> Allman, 1888		Hermanus, on beach: 34/19	
<i>H. inhacae</i> Millard, 1958		South of Sodwana: 27/32; coll. A. E. Louw	H 2891
<i>H. tenellum</i> Hincks, 1861		Hermanus, on beach: 34/19. Off Natal: 30/30 (vd); 31/30 (vd); MD	H 2934, 2938

APPENDIX 1 (continued)

Family and species	Locality data, including <i>S</i> —latitude/ <i>E</i> —longitude square	<i>SAM no.</i>
Lafoeidae		
<i>Acryptolaria conferta</i> (Allman, 1877)	Off Cape Town: 34/16 (md-d). Off Natal: 30/30 (vd); MD	H 1986, 2941
* <i>A. crassicaulis</i> (Allman, 1888)	Off Cape Town: 33/16 (a)	H 1988
<i>A. rectangularis</i> (Jarvis, 1922)	Off Natal: 30/30 (vd); MD	H 2929
<i>Cryptolarella abyssicola</i> (Allman, 1888)	Off Cape Town: 34/16 (md-d)	H 1987
<i>Filellum serratum</i> (Clarke, 1879)	Off Natal: 31/30 (vd); MD	H 2940
<i>Hebella scandens</i> (Bale, 1888)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	
<i>Lafoea dumosa</i> (Fleming, 1820)	Off Natal: 30/30 (vd); 31/30 (vd); MD	H 2944
<i>Zygophylax africana</i> Stechow, 1923	Off Natal: 30/30 (vd); MD	H 2930
<i>Z. sibogae</i> Billard, 1918	Off Natal: 30/30 (vd); MD	
Campanulariidae		
<i>Clytia gravieri</i> (Billard, 1904)	Sodwana Reef: 27/32 (s); coll. A. E. Louw. Off Cape Town: 34/17 (s); coll. Divn Sea Fisheries	H 2894, 2927
<i>Obelia dichotoma</i> (Linnaeus, 1758)	Hermanus, on beach: 34/19	
<i>O. geniculata</i> (Linnaeus, 1758)	Hermanus, on beach: 34/19	
Sertulariidae		
<i>Amphisbetia maplestonei</i> (Bale, 1884)	Jesser Point: 27/32 (l); coll. A. E. Louw. East London: 32/28; submitted by East London Museum	H 2895
<i>A. minima</i> (Thompson, 1879)	South of Sodwana: 27/32 (l); coll. A. E. Louw	H 2904
<i>Diphasia digitalis</i> (Busk, 1852)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2898
<i>D. tetraglochina</i> Billard, 1907	South of Sodwana: 27/32 (l); coll. A. E. Louw	H 2903
<i>Salacia articulata</i> (Pallas, 1766)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2888
<i>Sertularella diaphana</i> (Allman, 1886)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	
<i>S. fustiformis</i> (Hincks, 1861)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	
<i>S. leiocarpa</i> (Allman, 1888)	Stanford Bay: 34/19	H 2937
<i>S. mediterranea asymmetrica</i> Millard, 1958	Off Natal: 31/30 (vd); MD	H 2882
<i>S. m. mediterranea</i> Hartlaub, 1901	South of Sodwana: 27/32 (l); coll. A. E. Louw	
<i>Sertularia distans</i> (Lamouroux, 1816)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2896
<i>S. ligulata</i> Thornely, 1904	South of Sodwana: 27/32 (l); coll. A. E. Louw	H 2905
<i>S. longa</i> Millard, 1958	South of Sodwana: 27/32 (l); coll. A. E. Louw	H 2909
<i>Symplectoscyphus amphoriferus</i> (Allman, 1877)	Off Natal: 30/30 (vd); MD	H 2931

APPENDIX 1 (continued)

Family and species	Locality data, including S—latitude/E—longitude square	SAM no.
Plumulariidae		
<i>Aglaophenia pluma dichotoma</i> (M. Sars, 1857)	Sandwich Harbour: 23/14 (s); coll. B. Kensley	
<i>A. p. parvula</i> Bale, 1882	South of Sodwana: 27/32; coll. A. E. Louw	H 2942
<i>Antennella quadriaurita</i> Ritchie, 1909	Hermanus, on beach: 34/19	H 2901
<i>A. secundaria</i> (Gmelin, 1791)	South of Sodwana: 27/32; coll. A. E. Louw	H 2885
<i>Cladocarpus natalensis</i> Millard, 1977	Off Natal: 30/30 (vd); MD	
<i>Gattya multithecata</i> (Jarvis, 1922)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2932
<i>Gymnangium gracilicaule lignosum</i> (Millard, 1968)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2884
<i>Halopteris polymorpha</i> (Billard, 1913)	Off Natal: 30/30 (vd); MD	
<i>Lytocarpus filamentosus</i> (Lamarck, 1816)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2886
<i>L. phoeniceus</i> (Busk, 1852)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 309
<i>Monostachas natalensis</i> Millard, 1958	Agulhas Bank: 33/26 (d)	
<i>M. quadridens</i> (McCrary, 1858)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	
<i>Plumularia flicaulis</i> Kirchenpauer, 1876	South of Sodwana: 27/32 (l); coll. A. E. Louw. Near Lüderitz Bay: 26/14 (s)	H 2906, 2750
<i>P. obliqua</i> (Johnston, 1847)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2889
<i>P. pulchella</i> Bale, 1882	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2907
<i>P. setacea</i> (Linnaeus, 1758)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	
<i>Pycnotheca mirabilis</i> (Allman, 1883)	South of Sodwana: 27/32 (s); coll. A. E. Louw	H 2890
<i>Thecocarpus flexuosus flexuosus</i> (Lamouroux, 1816)	Sodwana Reef: 27/32 (s); coll. A. E. Louw	H 2887

<i>Gymnangium hians</i> (Busk, 1852)	Circum-tropical								X			X
<i>Halecium beanii</i> (Johnston, 1838)	Cosmopolitan	X		X	X	X	X	X	X	X	X	X
<i>Halecium delicatulum</i> Coughtrey, 1876	Cosmopolitan		X	X	X	X	X	X	X	X	X	X
<i>Halecium dichotomum</i> Allman, 1888	Endemic; W+S+E coast	X		X	X	X	X	X	X	X	X	X
<i>Halecium dyssymetrum</i> Billard, 1929	Circum-tropical								X			X
<i>Halecium halecinum</i> (Linnaeus, 1758)	Cosmopolitan					X			X			X
<i>Halecium inhacae</i> Millard, 1958	Endemic; E coast							X	X			
<i>Halecium lankesteri</i> (Bourne, 1890)	Scattered								X	X		X
<i>Halecium sessile</i> Norman, 1867	Cosmopolitan							X				X
<i>Halecium tenellum</i> Hincks, 1861	Cosmopolitan			X	X	X	X	X	X	X	X	X
<i>Halocordyle disticha</i> (Goldfuss, 1820)	Circum-tropical								X	X	X	X
<i>Halopteris gemellipara</i> Millard, 1962	Endemic; S + E coast			X	X	X	X		X			X
<i>Halopteris glutinosa</i> (Lamouroux, 1816)	Tropical; W.Ind.Oc.			X	X	X	X	X	X	X	X	X
<i>Halopteris polymorpha</i> (Billard, 1913)	Tropical; Ind.W.Pac.								X	X	X	X
<i>Halopteris pseudoconstricta</i> Millard, 1975	Endemic; W + S coast	X		X	X	X	X	X				X
<i>Halopteris rostrata</i> Millard, 1975	Endemic; E coast							X				X
<i>Halopteris tuba</i> (Kirchenpauer, 1876)	Endemic; S + E coast			X	X	X	X	X	X	X	X	X
<i>Hebella dispoliens</i> (Warren, 1909)	Endemic; E coast								X			
<i>Hebella furax</i> Millard, 1957	Tropical; W.Ind.Oc.			X	X	X	X	X	X	X	X	X
<i>Hebella parvula</i> (Hincks, 1853)	Temperate; Atlantic			X								X
<i>Hebella scandens</i> (Bale, 1888)	Cosmopolitan	X		X	X	X	X	X	X	X	X	X
<i>Hincksella corrugata</i> Millard, 1958	Tropical; W.Ind.Oc.							X	X	X	X	X

[illegible]

[illegible]



Millard, N A H. 1978. "The geographical distribution of southern African hydroids." *Annals of the South African Museum. Annale van die Suid-Afrikaanse Museum* 74, 159–200.

View This Item Online: <https://www.biodiversitylibrary.org/item/128445>

Permalink: <https://www.biodiversitylibrary.org/partpdf/92936>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.