

New Distributional Records for Two California Nudibranchs

BY

WILLIAM B. JAECKLE

207 Beryl Avenue, Mill Valley, California 94941

DURING SURVEYS OF INTERTIDAL opisthobranch mollusk communities in northern and central California, two nudibranchs were collected which represent a geographical range extension for each species. Both specimens were sighted in zone 3 rocky intertidal habitats (RICKETTS & CALVIN, 1968).

Ancula lentiginosa Farmer, in Farmer and Sloan, 1964

Relatively few sightings of the goniodorid *Ancula lentiginosa* have been reported since the original description. LANCE (1966) reported observing a specimen in a rocky intertidal region at Bahía de los Angeles, Baja California, establishing the southern extent of the geographical distribution. Along the California coast *A. lentiginosa* has been reported from La Jolla (FARMER & SLOAN, 1964), from Santa Barbara County (SPHON & LANCE, 1968), in San Luis Obispo County (ROLLER & LONG, 1969) and from Asilomar Beach, Monterey County (NYBAKKEN, 1978).

One specimen of *Ancula lentiginosa* was collected at the Frontier Arts Nature Reserve, Marin County (Lat. 37° 52' 22" N; Long. 122° 36' 56" W) on August 9, 1979. This specimen exhibited the solid freckled coloration pattern as described by FARMER & SLOAN (1964). Previous to this sighting, the northern limit of the geographical range was considered to be Elkhorn Slough, Monterey County (Gary McDonald, personal communication 5-XI-1979). The specimen collected in Marin County extends the northern limit of the geographical range 117.3 km. This specimen has been deposited in the invertebrate collection at the California Academy of Sciences, San Francisco, as a voucher specimen #1102.

Hallaxa chani Gosliner and Williams, 1975

Reported sightings of *Hallaxa chani* are limited primarily to the original description (GOSLINER & WILLIAMS, 1975). NYBAKKEN (1978) sighted a specimen at Asilomar Beach, Monterey County. McDONALD & NYBAKKEN (1978) reported 3 specimens from Morro Bay, San Luis Obispo County, feeding upon the encrusting ascidian *Didemnum carnulentum* Ritter and Forsyth, 1917.

A single specimen of *Hallaxa chani* was collected at Abalone Beach, Humboldt County (Lat. 41° 07' 20" N;

Long. 124° 09' 32" W) on April 28, 1979. The specimen was observed on the angiosperm *Phyllospadix scouleri* Hoof, 1839; GOSLINER & WILLIAMS (1975) cite *Phyllospadix* as an algal substrate particularly associated with *H. chani*. The external coloration and morphology of the collected specimen is consistent with the original description. The preserved specimen was sent to Dr. Terrence M. Gosliner who confirmed its identity.

The previous northernmost occurrence for this species is Bird Rock, Tomales Point, Marin County (Lat. 38° 14' N; Long. 123° 00' W); the southern extent of the geographical distribution is Shell Beach, San Luis Obispo County (Lat. 35° 12' N; Long. 120° 43' W) (GOSLINER & WILLIAMS, 1975). The collected specimen from Humboldt County represents a northern geographical range extension of 320 km.

ACKNOWLEDGMENTS

I would like to thank Dr. Gosliner for his support and encouragement and Mr. Gary McDonald for his assistance.

Literature Cited

- FARMER, WESLEY MERRILL & ALLAN J. SLOAN
1964. A new opisthobranch mollusk from La Jolla, California. *The Veliger* 6 (3): 148-150; pl. 18; 2 text figs. (1 January 1964)
- GOSLINER, TERRENCE M. & GARY C. WILLIAMS
1975. A genus of dorid nudibranch previously unrecorded from the Pacific coast of the Americas, with the description of a new species. *The Veliger* 17 (4): 396-405; 11 text figs. (1 April 1975)
- LANCE, JAMES ROBERT
1966. New distributional records of some northeastern Pacific Opisthobranchiata (Mollusca: Gastropoda) with descriptions of two new species. *The Veliger* 9 (1): 69-81; 12 text figs. (1 July 1966)
- McDONALD, GARY R. & JAMES WILLARD NYBAKKEN
1978. Additional notes on the food of some California nudibranchs with a summary of known food habits of California species. *The Veliger* 21 (1): 110-119 (1 July 1978)
- NYBAKKEN, JAMES WILLARD
1978. Abundance, diversity and temporal variability in a California intertidal nudibranch assemblage. *Mar. Biol.* 45 (2): 129-146; 15 text figs. (1 July 1977)
- RICKETTS, EDWARD F. & JACK CALVIN
1952. Between Pacific tides. Stanford Univ. Press, Stanford, Calif. v-xiii + 3-502; 46 pls.; 134 text figs.
- ROLLER, RICHARD A. & STEVEN J. LONG
1969. An annotated list of opisthobranchs from San Luis Obispo County, California. *The Veliger* 11 (4): 424-430 (1 April 1969)
- SPHON, GALE G. & JAMES ROBERT LANCE
1968. An annotated list of nudibranchs and their allies from Santa Barbara County, California. *Proc. Calif. Acad. Sci.* 36 (3): 73-84; 1 fig. (25 September 1968)



Jaekle, William B. 1981. "NEW DISTRIBUTIONAL RECORDS FOR 2 CALIFORNIA NUDIBRANCHS." *The veliger* 23, 240–240.

View This Item Online: <https://www.biodiversitylibrary.org/item/134165>

Permalink: <https://www.biodiversitylibrary.org/partpdf/93715>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.